arXiv:math/0109223v2 [math.AG] 22 Dec 2001

On Gorenstein log del Pezzo surfaces

Qiang Ye

Abstract

In this paper, we first present the complete list of the singularity types of the Picard number one Gorenstein log del Pezzo surface and the number of the isomorphism classes with the given singularity type. Then we give out a method to find out all singularity types of Gorenstein log del Pezzo surface. As an application, we present the complete list of the Dynkin type of the Picard number two Gorenstein log del Pezzo surfaces. Finally we present the complete list of the singularity type of the relatively minimal Gorenstein log del Pezzo surface and the number of the isomorphism classes with the given singularity type.

1 Introduction

Let V be a normal projective surface defined over the complex numbers field $\mathbf{C}$. We say that V is a *log del Pezzo surface* if V has at worst quotient singularities and if the anticanonical divisor $-K_V$ of V is an ample divisor. Then V is *Gorenstein*, i.e., K_V is a Cartier divisor, if and only if V has at worst rational double singularities. We note that a log del Pezzo surface is a rational surface (cf. [GZ]). The log del Pezzo surface has been studied from the various points of view. A general theory on the structure of such singular surfaces is developed in [M1] , [Z1] and [Z2]. The topological properties of the smooth part of the log del Pezzo surfaces and log Fano varieties are studied in [Fujiki], [GZ], [GZ1], [T1] and [Z3-Z6].

Let V be a Gorenstein log del Pezzo surface. Denote by $V^0 := V - (\mathrm{Sing}V)$ the smooth part of V. Let $g : U \longrightarrow V$ be a minimal resolution of singularities, we denote by $D := g^{-1}(\mathrm{Sing}V)$ the exceptional divisor and $\sharp(D)$ the number of the irreducible components of D.

In [Fu], the possible singularities of a Picard number one Gorenstein log del Pezzo surface are classified (see also [Brenton]).

Question 1.1 *Let V be a Picard number one Gorenstein log del Pezzo surface, is V then determined uniquely by its singularities?*

In general, V is not determined by its singularities. However, when V^0 is simply connected, this is almost the case. In [GPZ], [KM] and [MZ1], they have shown that for a Picard number one Gorenstein log del Pezzo surface V with simply connected smooth part, the singularity type of V is one of the following:

$$A_1,\ A_1+A_2,\ A_4,\ D_5,\ E_6,\ E_7,\ E_8.$$

Furthermore, V is uniquely determined by its singularities except in the case of $V(E_8)$ where there are exactly two isomorphism classes.

One of our main results is the following:

Theorem 1.2 *Let V be a Picard number one Gorenstein log del Pezzo surface ($\neq \mathbf{P}^2$), then the singularity types of V and the number m of the isomorphism classes of the Picard number one Gorenstein log del Pezzo surfaces with the given singularity type are listed in Table* 1.1.

Table 1.1

Dynkin type of V	m	*Dynkin type of V*	m	*Dynkin type of V*	m
E_8	2	$2A_4$	1	A_1+2A_3	1
A_1+E_7	2	$A_1+A_2+A_5$	1	E_6	1
D_8	1	$2A_1+2A_3$	1	A_1+A_5	1
A_8	1	$4A_2$	1	$3A_2$	1
A_2+E_6	2	E_7	1	D_5	1
$2A_1+D_6$	1	A_1+D_6	1	$2A_1+A_3$	1
$2D_4$	∞	A_7	1	A_4	1
A_3+D_5	1	A_2+A_5	1	A_1+A_2	1
A_1+A_7	1	$3A_1+D_4$	1	A_1	1

All isomorphism classes of any given singularity type are realizable (see the proof).

Corollary 1.3 *Let V be a Picard number one Gorenstein log del Pezzo surface. Then V is determined uniquely up to isomorphisms by the Dynkin type of V, except for $V(E_8)$, $V(A_1+E_7)$, $V(A_2+E_6)$ and $V(2D_4)$.*

Remark 1.4 *There are several results which are related to our Theorem* 1.2.

(1) *In* [Fu] *, the possible singularities of V are classified (see also* [Brenton] *).*

(2) *In* [MZ1]*, a classification of Picard number one Gorenstein log del Pezzo surfaces is given.*

In the proof, we need the result about the extremal rational elliptic surfaces which is proved in [MP] *but we do not use the classification mentioned in* [Brenton], [Fu] *and* [MZ1] *above.*

We also find out the singularity types of the Picard number two Gorenstein log del Pezzo surfaces.

Theorem 1.5 *Let V be a Picard number two Gorenstein log del Pezzo surface ($\neq \mathbf{P}^1 \times \mathbf{P}^1$ and Σ_1). Then there exist exactly* 45 *singularity types which are listed in Table* 1.2.

Table 1.2

E_7	A_7	$2A_3+A_1$	A_4+A_2	A_4+A_1	A_3
E_6+A_1	A_6+A_1	$3A_2+A_1$	$2A_3$	A_3+2A_1	A_2+A_1
D_7	A_5+A_2	E_6	$A_3+A_2+A_1$	$2A_2+A_1$	A_2
D_6+A_1	A_5+2A_1	D_6	A_3+3A_1	D_4	$2A_1$
D_5+A_2	A_4+A_3	D_5+A_1	$3A_2$	A_4	A_1
D_5+2A_1	$A_4+A_2+A_1$	D_4+2A_1	$6A_1$	A_3+A_1	
D_4+A_3	$A_3+A_2+2A_1$	A_6	D_5	A_2+2A_1	
D_4+3A_1	A_3+4A_1	A_5+A_1	A_5	$4A_1$	

All the above singularity types are realizable (see the proof).

The relatively minimal Gorenstein log del Pezzo surface (see Definition 2.1) is introduced in [MZ2] where the Dynkin type of the Picard number two relatively minimal Gorenstein log del Pezzo surface is given but the number of the isomorphism classes with the given singularity is not provided. In this paper, we prove the following theorem.

Theorem 1.6 *Let V be a Picard number two relatively minimal Gorenstein log del Pezzo surface ($\neq \mathbf{P}^1 \times \mathbf{P}^1$ and Σ_1). Then V has one of the* 10 *singularity types in Table* 1.3. *The number m of the isomorphism classes of the Picard number two relatively minimal Gorenstein log del Pezzo surfaces with the given singularity type are also listed in Table* 1.3.

Table 1.3

Dynkin type of V	m	*Dynkin type of V*	m	*Dynkin type of V*	m
D_7	∞	D_6	∞	D_4	1
D_5+2A_1	∞	D_4+2A_1	1	$4A_1$	1
A_3+4A_1	1	$2A_3$	∞		
A_3+D_4	∞	$6A_1$	1		

All isomorphism classes of the above singularity types are realized in the course of the proof.

Corollary 1.7 *Let* W *be a Gorenstein log del Pezzo surface.*

(1) *Then there are disjoint curves* $C_1, \cdots, C_k$ *and a birational morphism* $\sigma : W \longrightarrow W_{\min}$ *such that* $\sigma(C_i)$ *are smooth points ,* σ *is an isomorphism outside* $\cup C_i$ *and* $W_{\min}$ *is isomorphism to one of the surfaces* V *in Theorem* 1.2 *or* 1.6*.*

(2) *The singularity type of* W *is that of* V *plus a few type* A_{n_i} $(1 \leq i \leq k)$ *singularities lying on* C_i*.*

Remark 1.8 *From* [MZ2, Lemma 2] *or Lemma* 2.3*, we know that Theorems* 1.2 *and* 1.6 *present the complete list of the singularity types of the relatively minimal Gorenstein log del Pezzo surfaces and the number of the isomorphism classes with the given singularity type.*

This paper is organized as follows. In section 2, we introduce some terminology and preliminaries which will be used throughout the paper. In section 3, we establish the connection between Gorenstein log del Pezzo surfaces and rational elliptic surfaces. We will also give out a method to classify all singularity types of the Gorenstein log del Pezzo surfaces. In sections 4 and 5, we prove Theorems 1.2, 1.5 and 1.6.

Acknowledgement. I would like to thank Professor D.-Q. Zhang for many enlightening advice during the preparation of this paper.

2 Preliminaries

A. Relatively minimal Gorenstein log del Pezzo surfaces

Let V be a Gorenstein log del Pezzo surface. Let $f : U \longrightarrow V$ be a minimal resolution of singularities and let $D := f^{-1}(\mathrm{Sing}V)$. Then there are no $(-n)$-curves on U with $n \geq 2$ except for those (-2)-curves contained in D. The morphism f is the contraction of all (-2)-curves on U.

Let $\Sigma = \{F_1, \cdots, F_t\}$ be a maximum set of disjoint union of (-1)-curves on U such that the connected component of $\sum F_j + D$ containing F_i is a linear chain $F_i + \Delta_i$ of F_i and n_i $(n_i \geq 0)$ (-2)-curves with the dual graph:

$$(-1) - (-2) - (-2) - \cdots - (-2).$$

Let $U \longrightarrow U_{\min}$ be the smooth blow-down of all connected components of $\sum_{i=1}^{t}(F_i + \Delta_i)$. Let D' be the image on $U_{\min}$ of D and let $U_{\min} \longrightarrow V_{\min}$ be the contraction of D'. Then we have a commutative diagram:

$$\begin{tikzcd} U \arrow[r] \arrow[d] & U_{\min} \arrow[d] \\ V \arrow[r] & V_{\min} \end{tikzcd}$$

Here the map $V \longrightarrow V_{\min}$ contracts $\bar{F}_i := f(F_i)$ to t smooth points and is isomorphism outside $\cup \bar{F}_i$. Note that $\rho(V_{\min}) = \rho(V) - t$.

Definition 2.1 *The surface* $V_{\min}$ *is called a relatively minimal model of* V*.* V *is relatively minimal if* $V = V_{\min}$*. Note that* $V_{\min}$ *is not uniquely determined by* V*.*

From the above discussion, we have

Lemma 2.2 *Let* V *be a Gorenstein log del Pezzo surface,* U *the minimal resolution of* V *and* D *the exceptional divisor.*

(1) V *is not relatively minimal if and only if there is a* (-1)*-curve* E *on* U *such that the connected component of* $E + D$ *has dual graph:*

$$(-1) \quad \text{or} \quad (-1) - (-2) - (-2) - \cdots - (-2).$$

(2) *If* $\mathit{Sing}V_{\min}$ *has Dynkin type* $\sum a_i A_i + \sum d_i D_i + \sum e_i E_i$*, then* $\mathit{Sing}V$ *has Dynkin type* $\sum a_i A_i + \sum d_i D_i + \sum e_i E_i + \sum_{i=1}^{t} A_{n_i}$*.*

Thanks to the virtue of Miyanishi-Zhang (cf. [25, Lemma 2]), we have the following Lemma.

Lemma 2.3 *Let* V *be a Gorenstein log del Pezzo surface. Then there exists a birational morphism* $\eta : V \longrightarrow W$ *such that the following assertions hold.*

(1) W *is a relatively minimal Gorenstein log del Pezzo surfaces of Picard number one or two.*

(2) *Let* $E_1,\ldots,E_t$ *be all exceptional curves of* η*. Then* $w_i := \eta(E_i)(1 \leq i \leq t)$ *are* t *distinct smooth points of* W *for each* i*. Moreover* V *consists of the same singular points as on* W *and additional points* x_i *of the Dynkin type* A_{n_i} *with* $n_i \geq 1$*, where* $1 \leq i \leq t$*.*

B. Mordell-Weil groups of elliptic surfaces

(a) Definitions

Let C be a smooth projective curve over an algebraically closed field of characteristic 0, and Y a (Jacobian) *elliptic surface* over C. By this we mean the following: Y is a smooth projective surface with a relatively minimal elliptic fibration

$$f : Y \longrightarrow C$$

such that

(i) f has a global section $\mathcal{O}$, and

(ii) f is not smooth, i.e., there is at least one singular fibre.

For every (Jacobian) elliptic fibration Y there is a group structure of sections $MW(Y)$ with the distinguished section $\mathcal{O}$ as zero which we call *Mordell-Weil group.* Up to a finite group, $MW(Y)$ is identified with the relative automorphism group of the fibration. The basic results on the Mordell-Weil groups are proved in [Sh].

Due to a formula of Shioda-Tate we have the basic inequalities

$$0 \leq \mathrm{rank} MW(Y) \leq \rho(Y) - 2$$

where $\rho(Y)$ is the Picard number and the discrepancy in the upper bound is related to the degree of reducibility of the fibre.

Definition 2.4 *An elliptic fibration* Y *is called* extremal *if* $\rho(Y) = h^{1,1}(Y)$ (maximal Picard number) *and rank* $MW(Y) = 0$.

The extremal elliptic surfaces have been studied from various points of view. In [MP], the complete classification of extremal rational elliptic surfaces is given. For extremal elliptic $K3$ surfaces, we refer to [ATZ], [MP1], [MP2], [SZ] and [Ye].

Definition 2.5 Y *is called a* Mordell-Weil rank r *rational elliptic surface if* Y *is a rational surface and rank* $MW(Y) = r$.

In [M] and [P], the complete classification of the rational elliptic surfaces is given. In [OS], they describe the complete structure theorem of the Mordell-Weil groups of the rational elliptic surfaces.

The following result is proved in [OS, Corollary 2.3 and Theorem 2.5].

Theorem 2.6 *The Mordell-Weil rank one rational elliptic surface is generated by the section* P *which is disjoint from the zero-section* $\mathcal{O}$ *with* $< P, P > \leq 2$*, where* $< \cdot, \cdot >$ *is the height pairing on Mordel-Weil groups which is defined below.*

(b) Height Pairing

Let $f\colon Y \longrightarrow \mathbf{P}^1$ be a (relatively minimal) elliptic surface over $\mathbf{P}^1$ with a distinguished section $\mathcal{O}$. The complete list of possible fibres has been given by Kodaira [K1]. It encompasses two infinite families $(I_n, I_n^*, n \geq 0)$ and six exceptional cases $(II, III, IV, II^*, III^*, IV^*)$. And they can be considered as sublattices of the Néron-Severi group of Y.

Given an ellipric surface $f\colon Y \longrightarrow \mathbf{P}^1$, let $F_\nu = f^{-1}(\nu)$ denote the fibre over $\nu \in \mathbf{P}^1$, and let

$Sing(f) = \{\nu \in \mathbf{P}^1 | F_\nu \text{ is singular}\}$.

$\mathcal{R} = \mathrm{Red}(f) = \{\nu \in \mathbf{P}^1 | F_\nu \text{ is reducible}\}$.

For each $\nu \in \mathcal{R}$, let

$$F_\nu = f^{-1}(\nu) = \Theta_{\nu,0} + \sum_{i=1}^{m_\nu - 1} \mu_{\nu,i} \Theta_{\nu,i} \quad (\mu_{\nu,0} = 1)$$

where $\Theta_{\nu,i}$ $(0 \leq i \leq m_\nu - 1)$ are the irreducible components of F_ν, m_ν being their number, such that $\Theta_{\nu,0}$ is the unique component of F_ν meeting the zero section.

Lemma 2.7 (cf. [Sh]) *For each $\nu \in \mathbf{P}^1$, the intersection matrix*

$$A_\nu = ((\Theta_{\nu,i} \cdot \Theta_{\nu,j}))_{1 \le i,j \le m_\nu - 1}$$

is negative-definite.

Here we denote by $K = K(\mathbf{P}^1)$ the function field,

$$E(K) = \text{the group of sections of } \ f,$$

and

$$NS(Y) = \text{the group of divisors on } Y \ \text{ modulo algebraic equivalence.}$$

In general, for a rational elliptic surface Y with a section and with at least one singular fibre, there is a unique homomorphism φ of $E(K)$ into the Néron-Severi group $NS(Y) \otimes \mathbf{Q}$ such that $\varphi(P) \equiv (P) \bmod T_{\mathbf{Q}}$ and $\varphi(P) \perp T$ where T is the subgroup of $NS(Y)$ generrated by the zero section $\mathcal{O}$ and all the irreducible components of fibres (cf. [Sh, Lemma 8.1]).

Definition 2.8 *The pairing*

$$< P, Q >= -(\varphi(P) \cdot \varphi(Q))$$

on the Mordell-Weil group $E(K)$ is called the height pairing*, and the lattice*

$$(E(K)/E(K)_{tor}, <,>)$$

is called the Mordell-Weil Lattice *of the elliptic curve E/K or of the elliptic surface $f \colon Y \longrightarrow \mathbf{P}^1$.*

Theorem 2.9 (Explicit formula for the height pairing) [30, Theorem 8.6] *For any P,$Q \in E(K)$, we have*

$$< P, Q >= \chi + (P\mathcal{O}) + (Q\mathcal{O}) - (PQ) - \sum_{\nu \in R} contr_\nu(P, Q),$$

$$< P, P >= 2\chi + 2(P\mathcal{O}) - \sum_{\nu \in R} contr_\nu(P).$$

Remark 2.10 Here $\chi = \chi(\mathcal{O}_S)$, and $(P\mathcal{O})$ is the intersection number of the sections (P) and $(\mathcal{O})$, and similarly for $(Q\mathcal{O})$,(PQ). The term $contr_\nu(P, Q)$ stands for the local contribution at $\nu \in R$, which is defined as follows: suppose that (P) interests $\Theta_{\nu,i}$ and (Q) intersects $\Theta_{\nu,j}$. Then we let

$$contr_\nu(P, Q) = \begin{cases} (-A_\nu^{-1})_{i,j}, & if \ i \ge 1, j \ge 1, \\ 0, & otherwise. \end{cases}$$

where the first one means the (i, j)-entry of the matrix $(-A_\nu^{-1})$. Further we set

$$contr_\nu(P) = contr_\nu(P, P).$$

Arrange $\Theta_i = \Theta_{\nu,i}$ $(i = 0, 1, \cdots, m_\nu - 1)$ so that the simple components are numbered as in the figure below.



Figure 2.1

For the other types of reducible fibres, the numbering is irrelevant. Assume that (P) intersects $\Theta_{\nu,i}$ and (Q) intersect $\Theta_{\nu,j}$ with $i > 1$,$j > 1$. Then we have the following Table 2.1: the fourth row is for the case $i < j$ (interchange P, Q if necessary).

Table 2.1

T_ν^-	A_1	E_7	A_2	E_6	A_{b-1}	D_{b+4}
type of F_ν	III	III^*	IV	IV^*	$I_b(b \geq 2)$	$I_b^*(b \geq 0)$
$contr_\nu(P)$	$\frac{1}{2}$	$\frac{3}{2}$	$\frac{2}{3}$	$\frac{4}{3}$	$\frac{i(b-i)}{b}$	$\begin{cases} 1, & i = 1 \\ 1 + \frac{b}{4}, & i > 1 \end{cases}$
$contr_\nu(P, Q)(i < j)$	$-$	$-$	$\frac{1}{3}$	$\frac{2}{3}$	$\frac{i(b-j)}{b}$	$\begin{cases} \frac{1}{2}, & i = 1 \\ \frac{(2+b)}{4}, & i > 1 \end{cases}$

C. Weierstrass model of elliptic surfaces

Let $f : Y \longrightarrow C$ be a relatively minimal elliptic surface with the global section $\mathcal{O}$, and $K(Y)$ and $K(C)$ the function fields of Y and of C respectively. f induces a homomorphism $f^* : K(C) \longrightarrow K(Y)$, and $K(Y)$ is a transcendential extension of $K(C)$ of transcendence degree one. Y is birationaly equivalent to the subscheme Y^* in $Proj(\mathcal{O} \oplus \mathcal{O}(2L) \oplus \mathcal{O}(3L))$, which is given by the equation

$$V^2W = 4U^3 - g_2UW^2 - g_3W^3,$$

where $\mathcal{O}$ is the structure sheaf of C, L a line bundle on C with $degL = p_g - q + 1$, $g_2 \in H^0(C, \mathcal{O}(4L))$ and $g_3 \in H^0(C, \mathcal{O}(6L))$ are the sections with $\Delta = g_2^3 - 27g_3^2 \not\equiv 0$.

Theorem 2.11 (cf. [Kas]) *Y^* is an algebraic surface with rational double points as the only singularities. Y is the minimal resolution of Y^*. Y^* is determined by g_2, g_3 up to $\mathbf{C}^*$-operation*

$$(g_2, g_3) \longrightarrow (\lambda^4 g_2, \lambda^6 g_3), \lambda \in \mathbf{C}^* = \mathbf{C} - \{0\}.$$

g_2 , g_3 satisfy

(i) $\Delta = g_2^3 - g_3^2 \not\equiv 0$*,*

(ii) $\min(3\nu_p(g_2), 2\nu_p(g_3)) < 12$ *for all* $p \in C$*,*

where $v_p(g_2)$*,* $v_p(g_3)$ *and* $v_p(\Delta)$ *are the orders of the zeros of* g_2*,* g_3 *and* Δ *at* p*. The singular fiber in* Y^* *over* p *consists of the minimal resolution of the rational double point and the rational curve, which is defined by the section* $\mathcal{O}$*. The type of rational double and thereby the type of the singular fibre determines* $v_p(g_2)$*,* $v_p(g_3)$ *and* $v_p(\Delta)$*.* Y^* *is call the* Weierstrass model *of the elliptic surface.*

The $\mathcal{J}$*-invariant of the model is* $\mathcal{J} = \frac{g_2^3}{\Delta}$*.*

For the value of $v_p(g_2)$, $v_p(g_3)$ and $v_p(\Delta)$ of the singular fibre, we refer to [Her].

D. Classification of elliptic surfaces with at most 4 singular fibers

In [Hir], U.Schmickler-Hirzebruch classifies all relatively minimal elliptic fibrations Y over the complex projective line $\mathbf{P}^1$ which have at most three singular fibers. There are 36 possible combinations of singular fibers (each of the 36 potential representations yields different fiber types). Furthermore, U.Schmickler-Hirzebruch exhibits each of the 36 combinations of singular fibers on an elliptic surface of the desired type and shows that except for the trivial case (I_0, I_0, I_0) and the case (I_0^*, I_0^*), the surface is determined by the singular fiber types. On the other hand, except for the trivial case, those surfaces corresponding to rational or $K3$ surfaces. There are exactly *eleven* elliptic $K3$ surfaces with 3 singular fibers (cf. [SZ], [Ye]). Thus we get the following theorem.

Theorem 2.12 *Let* Y *be a rational elliptic surface with at least one and at most three singular fibers. Then except for singular fiber type* (I_0^*, I_0^*)*, the surface is unique up to isomorphisms.*

In [Her], S.Herfurtner studies the elliptic surfaces with four singular fibers and nonconstant $\mathcal{J}$-invariant over $\mathbf{P}^1$. He also distinguish two sets:

$$T^+ = \{I_n (n \geq 0), II, III, IV\} \text{ and } T^- = \{I_n^* (n \geq 0), IV^*, III^*, II^*\}.$$

Theorem 2.13 (cf. [Her]) *Let* $Y \longrightarrow \mathbf{P}^1$ *be a minimal elliptic surface with section, nonconstant* $\mathcal{J}$*-invariant and four singular fibers, of which at most one is in* T^-*.*

(i) If one singular fiber is in T^-*, three in* T^+*, the Weierstrass model depend on a parameter. Given four different base points, in the case* (I_1^*, I_1, I_1, II)*, there exists four, in the cases* (I_1^*, I_1, I_2, II) *and* (I_1, I_1, II, IV^*)*, there exist two elliptic surfaces, each depending on the* $\mathcal{J}$*-invariant, and for all other fiber combinations there exists precisely one elliptic surface.*

(ii) If all four singular fibers are in T^+*, then the Weierstrass models are determined uniquely up to isomorphisms, except for one combination:* (I_1, I_6, II, III)*, where there are two nonisomorphic models.*

E. Rational surfaces

Lemma 2.14 *Let Y be a smooth rational surface. Then we have*

(1) *Suppose that L is a smooth rational curve with $L^2 \geq 0$. Then $|L|$ is base point free and $h^0(Y, L) = 2 + L^2$.*

(2) *Suppose that L is an irreducible curve with $p_a(L) = 1$ and $L^2 \geq 1$. Then a general member of $|L|$ is smooth and $h^0(Y, L) = L^2 + 1$. If $L^2 \geq 2$, then $Bs|L| = \emptyset$. If $L^2 = 1$, then $|L|$ has exactly one base point.*

Proof. (1) is clear when $L^2 = 0$. So we assume $L^2 \geq 1$. Consider the exact sequence below

$$0 \longrightarrow \mathcal{O}_Y \longrightarrow \mathcal{O}_Y(L) \longrightarrow \mathcal{O}_{\mathbf{P}^1}(L^2) \longrightarrow 0.$$

Then, we have $h^0(Y, L) = 1 + h^0(\mathbf{P}^1, \mathcal{O}(L^2)) = 1 + (L^2 + 1) = L^2 + 2$.

To see $|L|$ is base point free, we let $\sigma : L_1 \longrightarrow L$ be the blow-up at $q \in L$ with E the exceptional curve and let L' be the proper transform of L. Then $L'^2 = L^2 - 1$ and $h^0(L') = L'^2 + 2 = L^2 + 1 = h^0(L) - 1$. On the other hand, $|L'| = |\sigma^*(L) - E| = |L - q|$; thus $|L|$ is base point free for

$$h^0(L - q) = h^0(L) - 1.$$

(2) Considering the similar exact sequence as in (1) and the equality

$$\chi(\mathcal{O}_L(L)) = h^0(L, \mathcal{O}_L(L)) - h^1(L, \mathcal{O}_L(L)) = \chi(\mathcal{O}_L) + deg\mathcal{O}_L(L),$$

we get

$$\begin{aligned} h^0(Y, L) &= 1 + h^0(L, \mathcal{O}_L(L)) = 1 + h^0(L, \omega_L(-L)) + 1 - p_a(L) + L^2 \\ &= 1 + L^2 + h^0(L, \omega_L(-L)) = 1 + L^2. \end{aligned}$$

Here ω_L is the dualizing sheaf and we have applied Serre duality for L. From this precise formula for $h^0(Y, L)$ and the similar discussion as in (1), we see that $Bs|L| = \emptyset$ if $L^2 \geq 2$. For the last assertion, see [DZ].

3 Reduction to the Rational Elliptic Surfaces

Before starting the results, we explain our terminology.

(a) Let V be a projective, normal surface with only rational double points as singularities. As usual, ratinal double points are indicated by their Dynkin types A_n, D_n, E_6, E_7 and E_8. When we say V a surface of type $2A_1 + D_6$ for example, it means that V has 3 singular points, one of which is of type D_6 and the other two are of type A_1. We indicate this by writing $V(2A_1 + D_6)$.

(b) For a log del Pezzo surface $V(2A_1+D_6)$ for example, we denote by $U(2A_1+D_6)$ the minimal resolution of $V(2A_1+D_6)$.

For readers' convenience, we give a short proof of a result in [5] using Kawamata-Viehweg Vanishing Theorem which was not available at that time.

Lemma 3.1 *Let U be the minimal resolution of a Gorenstein log del Pezzo surface V. Then $|-K_U|$ has a reduced irreducible member.*

Proof. Note that K_U is the pull back of K_V and $K_U^2 = K_V^2$ (cf. [1, 2]). Since $-K_U$, being the pull back of $-K_V$, is nef and big, by the Kawamata-Viehweg Vanishing theorem, we have $H^1(U, -K_U) = (0)$ (cf. [KMM]). Hence by the Riemann-Roch theorem, we get $\dim|-K_U| = K_U^2$. Suppose that a member A of $|-K_U|$ contains an arithmetic genus ≥ 1 irreducible component A_0. The Riemann-Roch theorem implies that $|A_0 + K_U| \neq \emptyset$. From this and $0 = A + K_U = (A_0 + K_U) + (A - A_0)$, we deduce that $A_0 = A$ with $p_a(A_0) = 1$ and Lemma 3.1 is true.

Thus we may assume that every member of $|-K_U|$ is a union of smooth rational curves. The Stein factorization and the fact that $q(U) = 0$ imply that a general member of $|-K_U|$ is of the form $M_1 + \cdots + M_k + F$, where F is the fixed part of the linear system, $M_i \cong \mathbf{P}^1$, and $M_i \sim M_j$.

Now suppose that $K_U^2 = 1$. If $F = 0$, then $k = 1$ and $M_1^2 = 1$ and Lemma 3.1 is true. Since $-K_U$ is nef and big, it is 1-connected by a result of C.P.Ramanujam (cf. [Reid]). Hence $1 = K_U^2 \geq (kM_1 + F).kM_1 \geq 1 + k^2M_1^2$. Thus $M_1^2 = 0$, $k = 1$, $M_1.F = 1$ and $K_U.F = 0$. Now intersecting the relation $-K_U \sim M_1 + F$ with the smooth rational curve M_1 of self-intersection 0, one gets a contraction. So Lemma 3.1 is true when $K_U^2 = 1$.

For $K_U^2 \geq 2$, let $U_1 \longrightarrow U$ be the blow-up of a point on $M_1 - (M_2 + \cdots + M_k + F)$. then $-K_{U_1}$ is linearly equivalent to the proper transform of $M_1 + \cdots + M_k + F$ and hence nef and big. If $U_1 \longrightarrow V_1$ is the contraction of all (-2)-curves then V_1 is a Gorenstein log del Pezzo surface. For U_1, we may argue as in the case of U and have reduced the proof of Lemma 3.1 to the case $K_U^2 = 1$, which has been dealt with in the previous paragraph. This completes the proof of Lemma 3.1.

Lemma 3.2 *Let U be the minimal resolution of a Picard number one Gorenstein log del Pezzo surface V with $K_V^2 = d$ where $2 \leq d \leq 7$. Then we can construct another surface U_{d-1} which is the minimal resolution of a Picard number one Gorenstein log del Pezzo surface V' with $K_{V'}^2 = d - 1$.*

Proof. From Lemma 3.1, we may choose an irreducible member of $|-K_U|$, say A and let E_d be a (-1)-curve on U. Such a (-1)-curve exists because $K_V^2 \leq 7$. Then A meets E_d at a point q. We may choose A general such that E_d is the only (-1)-curve through q. Let $U_{d-1} \longrightarrow U$ be the blow-up of q with E_{d-1} the except curve. Then $-K_{U_{d-1}}$ is linearly equivalent to the proper transform A_{d-1} of A. Since $K_{U_{d-1}}^2 > 0$, the divisor $-K_{U_{d-1}}$ is nef and big. And the curves having 0 intersection with A_{d-1} are precisely the inverse image of the (-2)-curves on U (contractible to singular points on V) and the proper transform E_d' of E_d. It follows that the contraction of all the (-2)-curves on U_{d-1} gives a Picard number one Gorenstein log del Pezzo surface V'.

Lemma 3.3 *Let V be a Picard number one Gorenstein log del Pezzo surface with $K_V^2 = 1$ and U the minimal resolution of V with D the exceptional divisor. Then there exists an extremal rational elliptic surface Y such that $g : Y \longrightarrow U$ is the blow-down of one (-1)-curve.*

Proof. Since K_U is the pull back of K_V, we have $K_U^2 = K_V^2 = 1$, $K_U^2 = 10-\rho(U)$, $\rho(U) = \rho(V) + \sharp(D)$ and $\sharp(D) = 8$. By Lemma 2.14, we know $Bs|-K_U| = \{p\}$.

Let $f : Y \longrightarrow U$ be the blow-up at p, then with Lemma 2.14, we have

(1) Y is a rational surface.

(2) $K_Y^2 = 0$ and $|-K_Y|$ induces an elliptic fibration of Y.

(3) $\rho(Y) = 10 - K_Y^2 = 10$.

(4) By the formula of Shioda (cf. [S])

$$10 = \rho(Y) = \sharp(D) + \text{rank MW(Y)} + 2,$$

we get rank MW(Y) = 0.

Thus Y is an extremal rational elliptic surface. This proves the lemma.

Combining Lemmas 3.2 and 3.3, we get the following Theorem 3.4.

Theorem 3.4 *Let V be a Picard number one Gorenstein log del Pezzo surface with $K_V^2 \leq 7$ and $U \longrightarrow V$ the minimal resolution . Then there exists an extremal rational elliptic surface Y such that $f : Y \longrightarrow U$ is a succession of blow downs of some (-1)-curves.*

We may generalize the above Theorem 3.4 to the Picard number r ($r \geq 2$) Gorenstein log del Pezzo surfaces.

Theorem 3.5 *Let V be a Picard number r ($r \geq 2$) singular Gorenstein log del Pezzo surface and $U \longrightarrow V$ the minimal resolution. Then there exists a Mordell-Weil rank $(r-1)$ rational elliptic surface Y such that $f : Y \longrightarrow U$ is a succession of blow downs of some (-1)-curves.*

The definition of the Mordell-Weil rank r rational elliptic surface is given in the Secition 2.

Proof. For simplicity, we only prove Theorem 3.5 for $r = 2$. The general case can be proved in the similar way.

We let $U \longrightarrow V$ be the minimal resolution of a Picard number two Gorenstein log del Pezzo surface V with D the exceptional divisor and $d := K_V^2 \geq 2$. From Lemma 3.1, we know that $|-K_U|$ has a reduced irreducible member.

Since $10 - K_V^2 = \rho(V) + \sharp(D)$, $\rho(V) = 2$ and $\sharp(D) \geq 1$ beacuse V is a singular surface, we get $10 - K_V^2 \geq 3$ and $K_V^2 \leq 7$. On the other hand, since K_U is the pull back of K_V, $K_U^2 = K_V^2 \leq 7$.

Thus we may choose an irreducible member of $|-K_U|$ say A and let E_d be a (-1)-curve on U. Such a (-1)-curve exists because $K_V^2 \leq 7$. Then A meets E_d at a point q.

Since $K_U^2 \geq 2$, we know $|-K_U|$ is base point free by Lemma 2.14. Furthermore since the smooth reduced irreducible member of $|-K_U|$ is dense in $|-K_U|$ and since there exists only finitely many (-1)-curves on U (cf. the proof of [6, Lemma 13]), we get the following Claim.

Claim 3.6 *We may choose* $A \in |-K_U|$ *such that there doesn't exist another* (-1) *curve passing through the point* q.

Let U_{d-1} be the blow-up of q with E_{d-1} the exceptional curve and E'_d the proper transform of E_d. Then by the similar discussion as Lemma 3.2, we know that U_{d-1} is the minimal resolution of a Gorenstein log del Pezzo surface V' with $K_{V'}^2 = d-1$.

Claim 3.7 $\rho(V') = 2$.

Proof of the Claim. It is clearly that $\rho(V') \leq 2$. Thus to prove the Claim, we need to show that $\rho(V') \neq 1$. Suppose $\rho(V') = 1$. Then from the above construction, we know that there must exist a curve C on U such that $C^2 \geq 0$ and $mult_p C \geq 2$ and the proper transform C' of C in U_{d-1} is an (-2)-curve. Thus we have $C'.E_{d-1} = mult_p C = m \geq 2$, $E'_d.E_{d-1} = 1$ where C' and E'_d are (-2)-curves.

Now we pay attention to the set $I = \{E'_d, E_{d-1}, C'\}$ in U_{d-1}. Clearly we may do the same operation (blow-up) on U_{d-1} as before and finally get an rational elliptic surface Y. Since C' and E'_d are already (-2)-curves we will keep the same symbols in Y and let E' be the proper transform of E_{d-1}. To prove the Claim, we need to consider two cases.

1. E' is an (-1)-curve.

In this case and from the elliptic fibration of Y, we see that C' and E'_d will be in two different elliptic fibers and E' is a section of Y which is a contradiction since $C'.E' = C'.E_{d-1} \geq 2$.

2. E' is an (-2)-curve.

In this case we see that E'_d, E' and C' will be in one elliptic fiber. We also get a contracdiction since $C'.E' \geq 2$ and we can't find an elliptic fiber contains C' and E'.

Thus we prove the Claim. We note that the above proof can easily be generalized to prove the general case.

Continue this process. Since $K_{U_{d-1}}^2 = d-1 < K_U^2$, this process must stop at the $(d-1)$-th step. Thus, we obtain a surface U_1 which is the minimal resolution of the Picard number two Gorenstein log del Pezzo surface V_1 with $K_{V_1}^2 = 1$. On U_1 we can not do the above process again. But since $K_{U_1}^2 = 1$, $|-K_{U_1}|$ has exactly one base point q by Lemma 2.14. Blowing up q will give rise to a Mordell-Weil rank one rational elliptic surface Y as in Lemma 3.3. Thus Theorem 3.5 is proved.

Remark 3.8 *From Theorems* 3.4 *and* 3.5*, together with* [OS] *and* [P]*, we may classify all singularity types of the Gorenstein log del Pezzo surfaces.*

Question 3.9 *From the proofs of Theorems* 3.4 *and* 3.5*, it is natural for us to ask the converse question: If* $U \longrightarrow V$ *is the minimal resolution of a Picard number one (resp. Picard number* $r \geq 2$*) Gorenstein log del Pezzo surface* V *. Let* E *be a* (-1)*-curve on* U *and* $\phi : U \longrightarrow U'$ *the blow-down of* E*, then could* U' *be a minimal resoultion of a Picard number one (resp. Picard number* $r \geq 2$*) Gorenstein log del Pezzo surface* V'*?*

Lemma 3.10 *Let the notation be the same as in Question* 3.9*. Let* $f : U \longrightarrow U'$ *be the blow-down of a* (-1)*-curve* E *and* D *the exceptional divisor of the minimal resolution* $U \longrightarrow V$*. Let* $U' \longrightarrow V'$ *be the contraction of all* (-2)*-curves and let* D' *be the exceptional divisor. Then*

(1) V' *is a Gorenstein log del Pezzo surface.*

(2) *If* $D.E = 1$*,* $\rho(V') = \rho(V)$*. The converse is also true.*

Proof. We fix a nonsingular member $A \in |-K_U|$, the existence of which is proven in [D] . We also let $A' = f_*(A)$. Then $A' \in |-K_{U'}|$ and $(A'^2) \geq (A^2) \geq 1$. Since $-K_U$ is nef and big, so is $-K_{U'}$. To vertify (1), we have only to show that $-K_{V'}$ is ample. In other words, for every curve C on U' with $(-K_{U'}, C) = 0$, we must show that $C \leq D'$. Since $-K_{U'}$ is nef and big, by Hodge Index Theorem, we get $C^2 < 0$. On the other hand,

$$2p_a(C) - 2 = C^2 + C.K_{U'} = C^2 < 0.$$

Thus we have $p_a(C) = 0$ and so $C = \mathbf{P}^1$, $C^2 = -2$ and $C \leq D'$. (1) is proved.

(2) Since $E.D = 1$, we have $K_{U'}^2 = K_U^2 + 1$ and $\sharp(D) = \sharp(D') + 1$. Thus the first part in the assertion (2) follows from the relations $K_U^2 = K_V^2 = 10 - \rho(U)$, $K_{U'}^2 = K_{V'}^2 = 10 - \rho(U')$, $\rho(U) = \rho(V) + \sharp(D)$and $\rho(U') = \rho(V') + \sharp(D')$. The converse part is clear from the above discussion.

We will call a (-1)-curve which satisfies the part (2) of Lemma 3.10 a *Nice Exceptional Curve* or simply an *NEC*.

4 Complete Classification of the Picard number one Gorenstein log del Pezzo Surfaces

We shall prove Theorem 1.2 in the present section.

Let V be a Picard number one Gorenstein log del Pezzo surfaces and $g : U \longrightarrow V$ a minimal resolution of singularities. Then $K_U^2 = K_V^2 > 0$, $|-nK_U| \neq \phi$ for $n > 0$ and U is a rational surface. Hence $K_U^2 + \rho(U) = 10$. This implies that $1 \leq K_U^2 \leq 9$. We observe:

(i) If $K_U^2 = 9$, then V is smooth and isomorphic to $\mathbf{P}^2$.

(ii) If $K_U^2 = 8$, then U is the Hirzebruch surface Σ_2 and V is obtained by contracting the (-2)-curve to an A_1-singularity. In this case V is isomorphic to the quadric cone $Q := \{X^2 + Y^2 + Z^2 = 0\}$ in $\mathbf{P}^3$.

Lemma 4.1 *$K_U^2 \neq 7$.*

Proof. Assume the contrary, i.e., $K_U^2 = 7$. In this case U is obtained from a Hirzebruch surface Σ_n by one blow-up with E the exceptional curve. Let S be a curve on Σ_n with self-intersection $-n$ and L a fibre of the $\mathbf{P}^1$-fibration on Σ_n. We may assume that the blow-up point $p \in L$. We have $K_{\Sigma_n} \sim -2S - (2+n)L$.

If $p \notin S$, then $K_U \sim -2S - (2+n)L' - (1+n)E$, where L' is the proper transform of L. Clearly, U contains a (-2)-curve different from S, say C. Since $K_U.C = 0$, the curve C is disjoint from S, L' and E. This is impossible.

If $p \in S$, then $K_U \sim -2S' - (2+n)L' - (3+n)E$ where S' is the proper transform of S. Again there is a (-2)-curve different from S. Thus we get a contradiction as above. This completes the proof of the lemma.

In the rest of this section, we assume $1 \leq K_V^2 \leq 6$.

We start with $K_V^2 = 1$.

From Lemma 3.3, we know that if U is the minimal resolution of a Picard number one Gorenstein log del Pezzo surface V, then every NEC on U comes from either the (-1)-curve on an extremal rational elliptic surface Y or the (-2)-curve which intersects with the exceptional curve of the blow-down $Y \longrightarrow U$. The (-1)-curves on Y are the sections of the elliptic fibration and form the Mordell-Weil group of the elliptic fibration $Y \longrightarrow \mathbf{P}^1$.

From [MP], we get a complete list of the extremal rational elliptic surfaces Y which are listed in Table 4.1. For convenience, we label the surfaces. By contracting the zero section $\mathcal{O}$ in $MW(Y)$, we get a surface U which is the minimal resolution of a Picard number one Gorenstein log del Pezzo surface V with $K_V^2 = 1$. By a surface U_j^i, we mean that it is the blow-down of some (-1)-curves from the No.i extremal rational elliptic surface and $K^2 = j$. We let n be the number of the NEC on U_1.

Table 4.1

♯	Y	U_1	n	♯	Y	U_1	n
1	II, II^*	$U_1^1(E_8)$	1	9	I_1^*, I_4, I_1	$U_1^9(D_5 + A_3)$	1
2	III, III^*	$U_1^2(A_1 + E_7)$	1	10	I_2^*, I_2, I_2	$U_1^{10}(D_6 + 2A_1)$	1
3	IV, IV^*	$U_1^3(A_2 + E_6)$	1	11	I_9, I_1, I_1, I_1	$U_1^{11}(A_8)$	2
4	I_0^*, I_0^*	$U_1^4(2D_4)$	2	12	I_8, I_2, I_1, I_1	$U_1^{12}(A_7 + A_1)$	1
5	II^*, I_1, I_1	$U_1^5(E_8)$	1	13	I_5, I_5, I_1, I_1	$U_1^{13}(2A_4)$	0
6	III^*, I_2, I_1	$U_1^6(E_7 + A_1)$	1	14	I_6, I_3, I_2, I_1	$U_1^{14}(A_5 + A_2 + A_1)$	0
7	IV^*, I_3, I_1	$U_1^7(E_6 + A_2)$	1	15	I_4, I_4, I_2, I_2	$U_1^{15}(2A_1 + 2A_3)$	0
8	I_4^*, I_1, I_1	$U_1^8(D_8)$	2	16	I_3, I_3, I_3, I_3	$U_1^{16}(4A_2)$	0

From the proof of Lemma 3.3, we know there is a one to one correspondence between the No.i extremal rational elliptic surfaces Y and U_1^i. On the other hand, for every $i \neq 4$, there is only one isomorphism class of the No.i extremal rational elliptic surface (cf. [MP, Theorem 5.4]). The extremal rational elliptic surfaces of No.4, i.e., with singular fibers (I_0^*, I_0^*), are parameterized by $j \in \mathbf{C}$ [ibid.]. Thus we get the following lemma:

Lemma 4.2 *Let V be the Picard number one Gorenstein log del Pezzo surface with $K_V^2 = 1$. Then the singularity type of V, the number m of the isomorphism class with the given singularity type are listed in Table 4.2.*

Table 4.2

Dynkin type of V	m	*Dynkin type of* V	m
E_8	2	A_8	1
$A_1 + E_7$	2	$A_1 + A_7$	1
$A_2 + E_6$	2	$2A_4$	1
$2D_4$	∞	$A_1 + A_2 + A_5$	1
D_8	1	$2A_1 + 2A_3$	1
$A_3 + D_5$	1	$4A_2$	1
$2A_1 + D_6$	1		

Next we consider $K_V^2 = 2$.

By blowing down an NEC on U_1 which is listed in Table 4.3, we get a surface U_2 which is the minimal resolution of a Picard number one Gorenstein log del Pezzo surface V_2. They are listed in the following table where we let n be number of the NECs on U_2.

Table 4.3

U_1	U_2	n	U_1	U_2	n
$U_1^1(E_8)$	$U_2^1(E_7)$	1	$U_1^8(D_8)$	$U_2^8(A_1 + D_6)$	1
$U_1^2(A_1 + E_7)$	$U_2^2(A_1 + D_6)$	1	$U_1^8(D_8)$	$U_2^8(A_7)$	2
$U_1^3(A_2 + E_6)$	$U_2^3(A_2 + E_5)$	1	$U_1^9(A_3 + D_5)$	$U_2^9(A_1 + 2A_3)$	0
$U_1^4(2D_4)$	$U_2^4(3A_1 + D_4)$	0	$U_1^{10}(2A_1 + D_6)$	$U_2^{10}(3A_1 + D_4)$	0
$U_1^5(E_8)$	$U_2^5(E_7)$	1	$U_1^{11}(A_8)$	$U_2^{11}(A_2 + A_5)$	1
$U_1^6(A_1 + E_7)$	$U_2^6(A_1 + D_6)$	1	$U_1^{12}(A_1 + A_7)$	$U_2^{12}(A_1 + 2A_3)$	0
$U_1^7(A_2 + E_6)$	$U_2^7(A_2 + A_5)$	1			

Remark 4.3 (i) *In the following discussion, the Figure i ($i = 1, 2, 3, 4$) are listed in the configurations of Appendix.*

*(ii) To see $U_2^{11}(A_2 + A_5)$ also doesn't depend on the choice of the blow-down curve $E_i (i = 1, 2)$ on $U_1^{11}(A_8)$ (cf. Figure 2), we look at the No.*11 *extremal rational elliptic*

surface Y (cf. Table 2). A simple calculation shows that the Mordell-Weil group of Y is $\mathbf{Z}/3\mathbf{Z}$ and there are three sections E_0,E_1 and E_2 (cf. Figure 2). The section E_1 defines an automorphism g of Y such that $g(E_0) = E_1$, $g(E_1) = E_2$ and $g(E_2) = E_0$. For simplicity, we denote by bd_i the blow-down of the E_i ($i = 0, 1, 2$). Then

$$bd_2(U_1^{11}(A_8)) = bd_2 \circ bd_0(Y) \cong bd_0 \circ bd_1(Y) \cong bd_1 \circ bd_0(Y) = bd_1(U_1^{11}(A_8)).$$

Claim 4.4 $U_2^1(E_7) \cong U_2^5(E_7)$.

Proof. For simplicity, we let

$$\begin{array}{llll} U_1 = U_1^1(E_8), & V_1 = V_1^1(E_8), & U_2 = U_2^1(E_7), & V_2 = V_2^1(E_7), \\ U_1' = U_1^5(E_8), & V_1' = V_1^5(E_8), & U_2' = U_2^5(E_7), & V_2' = V_2^5(E_7). \end{array}$$

By the construction of U_2, U_2', there is a unique (-1)-curve E_2 on U_2. To prove $U_2 \cong U_2'$, we only need to show that $|-K_{U_2}|$ has a cuspidal member A which is the image of the type II fiber and also a nodal member A'; in fact if $U_1 \to U_2$ (resp. $U_1' \to U_2$) is the blow-up of $q = A \cap E_2$ (resp. $q' = A' \cap E_2$) then $|-K_{U_2}|$ (resp.$|-K_{U_2'}|$) has a cuspidal (resp. nodal) member, whence

$$U_2 \cong \text{ (the blow − down of the unique (−1) − curve on U}_1' \text{) } = U_2'.$$

We let Y (resp. Y') be the extremal rational elliptic surface with singular fiber type II^*, II (resp. II^*,I_1,I_1). There is a composition $Y \longrightarrow \mathbf{P}^2$ (resp. $Y' \longrightarrow \mathbf{P}^2$) of blow-down of the section E and all components in the type II^* fiber except for a multiplicity-3 component C_3'.

Clearly the above birational morphism $Y \longrightarrow \mathbf{P}^2$ factors as $Y \longrightarrow U_2 \longrightarrow \mathbf{P}^2$, where $Y \longrightarrow U_2$ is the blow-down of E and the multiplicity-1 component C_1 of II^*.

Thus the existence of A, A' on U_2 is equivalent to the existence of a cuspidal cubic B (the image of the type II fibre) and a nodal cubic B' on $\mathbf{P}^2$ with a common inflex (the image of the multiplicity-3 component C_3' in the type II^* fibre) defined by $Z = 0$, with local intersection $I(B, B'; [0, 1, 0]) = 7$ at the inflex point $[0, 1, 0]$. We may assume that $B = \{Y^2Z = X^3\}$ and just let $B' = \{Y^2Z = X^3 + XZ^2 + \sqrt{-4/27}\}$. This completes the proof of the Claim.

Claim 4.5 (1) $U_2^2(A_1 + D_6) \cong U_2^6(A_1 + D_6) \cong U_2^8(A_1 + D_6)$ (cf. Figure 1).
(2) $U_2^3(A_2 + A_5) \cong U_2^7(A_2 + A_5) \cong U_2^{11}(A_2 + A_5)$ (cf. Figure 2).
(3) $U_2^4(3A_1 + D_4) \cong U_2^{10}(3A_1 + D_4)$ (cf. Figure 3).
(4) $U_2^9(A_1 + 2A_3) \cong U_2^{12}(A_1 + 2A_3)$ (cf. Figure 4).

Proof. (1) From the construction of $U_2^2(A_1 + D_6)$, there is an NEC E_2 on $U_2^2(A_1 + D_6)$ and there is a smooth rational curve L and $L^2 = 0$ such that $L \cap E_2 = \{p\}$. On the other hand, by Lemma 2.14, we know that $|L|$ is base point free and defines a $\mathbf{P}^1$-fibration. We choose a general member L_1 of $|L|$, and let $\{q\} = L_1 \cap E_2$. It is easy to see that $U_1^2(A_1+E_7)$ is the blow-up of $U_2^2(A_1+D_6)$ at point p and $U_1^6(A_1+E_7)$ is the blow-up of $U_2^2(A_1+D_6)$ at point q. Thus

$$U_2^2(A_1+D_6) \cong (\text{ the blow}-\text{down of the uniqe NEC on } U_1^6(A_1+E_7)) = U_2^6(A_1+D_6)$$

On the other hand, blowing up at point r in $U_2^6(A_1+D_6)$ will produce $U_1^8(D_8)$, and there is unique way from $U_1^8(D_8)$ to $U_2^8(A_1+D_6)$. Thus

$$U_2^2(A_1+D_6) \cong U_2^8(A_1+D_6).$$

By the similar argument as (1), we may prove (2), (4).

(3) Since $U_1^{10}(2A_1+D_6)$ is unique and there is only one NEC on $U_1^{10}(2A_1+D_6)$, we know $U_2^4(3A_1+D_4)$ is unique and

$$U_2^4(3A_1+D_4) \cong U_2^{10}(3A_1+D_4).$$

This proves the Claim.

Since $U_i^j \cong U_{i'}^{j'}$ if and only if $V_i^j \cong V_{i'}^{j'}$, we get the following lemma:

Lemma 4.6 *Let V be the Picard number one Gorenstein log del Pezzo surface with $K_V^2 = 2$. Then the singularity type of V is one of the following:*

$$E_7, A_7, A_1+D_6, A_1+2A_3, A_2+A_5, 3A_1+D_4.$$

Furthermore, V is uniquely determined by its singularity type.

Next we deal with $K_V^2 = 3$.

By blowing down an NEC on U_2 which is listed in Table 4.4, we get a surface U_3 which is the minimal resolution of a Picard number one Gorenstein log del Pezzo surface V_3. They are listed in the following table where we let n be the number of the NEC on U_3.

Table 4.4

U_2	U_3	n	U_2	U_3	n
$U_2^1(E_7)$	$U_3^1(E_6)$	1	$U_2^3(A_2+A_5)$	$U_3^3(3A_2)$	0
$U_2^6(A_1+D_6)$	$U_3^6(A_1+A_5)$	1	$U_2^8(A_7)$	$U_3^8(A_1+A_5)$	1

Remark 4.7 (*i*) *There are two NECs, E_0' and E_1' on $U_2^8(A_7)$ (cf. Figure 1). To see $U_3^8(A_1+A_5)$ doesn't depend on the choice of the blow-down of the E_i' on $U_2^8(A_7)$ where $i = 1, 2$, we look at the No.8 extremal rational elliptic surface Y (cf. Table 2). A direct calculation shows that the Mordell-Weil group of Y is $\mathbf{Z}/2\mathbf{Z}$ and there are two sections E_0, E_1 (cf. Figure 1). The section E_1 define an automorphism g of Y which interchanges E_0 and E_1. And this g will induce an automorphism of $U_2^8(A_7)$*

which interchanges E_i' $(i = 0, 1)$*. Thus blowing down* E_0 *or* E_1 *will produce the same surface.*

(ii) *By the similar argument as in Claim* 4.5*, we have*

$$U_3^8(A_1 + A_5) \cong U_3^6(A_1 + A_5) \quad \text{(cf.Figure 1).}$$

Thus we have the following lemma:

Lemma 4.8 *Let* V *be the Picard number one Gorenstein log del Pezzo surface with* $K_V^2 = 3$*. Then the singularity type of* V *is one of the following:*

$$E_6, 3A_2, A_1 + A_5.$$

V *is uniquely determined by its singularity type.*

Finally we deal with $4 \leq K_V^2 \leq 6$.

By blowing down an NEC on U_3 which is listed in Table 4.5, we get a surface U_4 which is the minimal resolution of a Picard number one Gorenstein log del Pezzo surface V_4.

Table 4.5

U_3	U_4	n
$U_3^1(E_6)$	$U_4^1(D_5)$	1
$U_3^8(A_1 + A_5)$	$U_4^8(2A_1 + A_3)$	0

Lemma 4.9 *Let* V *be the Picard number one Gorenstein log del Pezzo surface with* $K_V^2 = 4$*. Then the singularity type of* V *is one of the following:*

$$D_5, 2A_1 + A_3$$

Furthermore, V *is uniquely determined by its singularity type.*

By blowing down the unique NEC on $U_4^1(D_5)$, we get $U_5^1(A_4)$ and so the Picard number one Gorenstein log del Pezzo surface $V(A_4)$. Then blowing down the unique NEC on $U_5^1(A_4)$, we get $U(A_1 + A_2)$ with $\sharp\{NEC \text{ on } U(A_1 + A_2)\} = 0$, and so the Picard number one Gorenstein log del Pezzo surface $V(A_1 + A_2)$.

Combining Lemmas 4.2, 4.6, 4.8, 4.9 and the above discussion, we prove Theorem 1.2.

5 Complete classification of the Picard number two relatively minimal Gorenstein log del Pezzo Surfaces

We shall prove Theorems 1.5 and 1.6 in the present section.

Let V be a Picard number two Gorenstein log del Pezzo surface and $g : U \longrightarrow V$ a minimal resolution of singularities. We also denote by $D := g^{-1}(\mathrm{Sing}V)$ the exceptional divisor.

We start with $K_V^2 = 1$.

From Theorem 3.5,we know that any rational surface U which is the minimal resolution of a Picard number two Gorenstein log del Pezzo surface V with $K_V^2 = 1$ comes from the Mordell-Weil *rank one* rational elliptic surface Y. Meanwhile, every NEC on U comes from either the (-1)-curve on Y or the (-2)-curve which intersects the NEC of the blow-down $Y \longrightarrow U$. The (-1)-curves on Y are the sections of the elliptic fibration and form the Mordell-Weil group of the elliptic fibration $Y \longrightarrow \mathbf{P}^1$.

From [M], [OS] and [P], we get a complete list of the Mordell-Weil *rank one* rational elliptic surfaces Y and its Mordell-Weil group. By contracting the zero section $\mathcal{O}$ in $MW(Y)$, we get a surface U_1 which is the minimal resolution of a Picard number two Gorenstein log del Pezzo surfaces V with $K_V^2 = 1$. For convenience, we label the surfaces which are listed in Table 1 of Appendix. By a surface U_j^i, we mean that it is the smooth blow-down of some (-1)-curves from the No.i Mordell-Weil rank one rational elliptic surface and $K^2 = j$. Thus we get the following lemma.

Lemma 5.1 *Let V be the Picard number two Gorenstein log del Pezzo surface with $K_V^2 = 1$. Then there exist* 18 *singularity types which are listed in Table* 5.1.

Table 5.1

E_7	$D_5 + A_2$	A_7	$A_4 + A_3$	$2A_3 + A_1$
$E_6 + A_1$	$D_5 + 2A_1$	$A_1 + A_6$	$A_4 + A_2 + A_1$	$3A_2 + A_1$
D_7	$D_4 + A_3$	$A_5 + A_2$	$A_3 + A_2 + 2A_1$	
$D_6 + A_1$	$D_4 + 3A_1$	$A_5 + 2A_1$	$A_3 + 4A_1$	

In order to find out the list of all Picard number two *relatively minimal* Gorenstein log del Pezzo surfaces, we need to exclude the surfaces on which there is a (-1)-curve E and a connected component R of D (R may be zero) such that $E + R$ is a linear chain and $E + R$ is a connected component of $E + D$ (cf. Table 1 of Appendix). In fact we have the following lemma:

Lemma 5.2 *None of U_1^i ($7 \leq i \leq 38$) is the minimal resolution of a Picard number two relatively minimal Gorenstein log del Pezzo surface.*

Proof. We verify the above assertions by checking each case separately. To exhibit our arguments, we treat only No.12 and No.24, i.e., $U_1^{12}(2A_3+A_1)$ and $U_1^{24}(3A_2+A_1)$ respectively. The other cases can be treated similarly. We also write out the linear chain exists in the corresponging surface in Table 1 of the Appedenix.

(1) $U_1^{12}(2A_3+A_1)$.

The Mordell-Weil group of the No.12 Mordell-Weil rank one rational elliptic surface Y (with fibre type $2I_4, I_2, 2I_1$) is $A_1^* \oplus \mathbf{Z}/4\mathbf{Z}$ where A_1^* is the rank one lattice generated by an element e such that $< e, e >= 1/2$. We let P be the generator of the lattice A_1^* and Q the generator of the torsion part of the Mordell-Weil group,i.e., $\mathbf{Z}/4\mathbf{Z}$. A direct calculation shows that there is one element in the set $\{P_2, P_2+Q, P_2+2Q, P_2+3Q\}$ where $P_2 := 2P$, say P_2, passing through the 0-th components of the singular fibres $2I_4, I_2$ and $(P_2\mathcal{O}) = 0$. Since $U_1^{12}(2A_3+A_1)$ is the blow-down of the zero section $\mathcal{O}$ of Y, the image of P_2 will be the (-1)-curve disjoint from $D = 2A_3+A_1$. Thus $U_1^{12}(2A_3+A_1)$ is not the minimal resolution of a Picard number two relatively minimal Gorenstein log del Pezzo surface.

(2) $U_1^{24}(3A_2+A_1)$.

To start with, we let P be the generator of the rank 1 lattice $< 1/6 >$. Firstly we need to find out all the (-1)-curves on $U_1^{24}(3A_2+A_1)$. Since we blow down the zero section $\mathcal{O}$ on the No.24 Mordell-Weil rank one rational elliptic surface with the fibre type $(3I_3, I_2, I_1)$, the section $P_m := mP$ $(m \in \mathbf{Z})$ which intersects $\mathcal{O}$ will not be a (-1)-curves on $U_1^{24}(3A_2+A_1)$ again. Thus we get the following claim:

Claim 5.3 *A (-1)-curve on U_1 is either the image of the zero component of the singular fibre or the image of the section disjoint from the $\mathcal{O}$.*

Suppose $(P\mathcal{O}) = 0$. Then from $m^2/6 =< P_m, P_m >= 2\chi+2(P_m\mathcal{O})-\sum_{\nu\in R} contr_\nu(P_m) = 2-\sum_{\nu\in R} contr_\nu(P_m) \leq 2$, we get $m^2 \leq 12$ and $|m| \leq 3$. Conversely, a simple calculation shows that $(P_3\mathcal{O}) = 0$ and P_3 only passes through the 1-st component of I_2. Thus P_3+A_1 will form a linear chain required.

By a similar discussion as in Lemma 4.2, we know there is a one to one correspondence between No.i Mordell-Weil rank one rational elliptic surfaces and U_1^i, where $1 \leq i \leq 6$. Thus finding the number of the isomorphism classes of the relatively minimal Dynkin type on $K_V^2 = 1$ is equivalent to finding the number of the isomorphism classes of the rational elliptic surfaces with the corresponding singular fibre type.

Remark 5.4 (1) *From* [Hir] *or Theorem* 2.12 *, we know the Nos.*1 *and* 4 *Mordell-Weil rank one rational elliptic surfaces are unique up to isomorphism,i.e., are determined by the singular fibre types.*

(2) *From* [Her] *or Theorem* 2.13*, we know that the number of the isomorphism classes of each of the Nos.* 2*,* 5 *and* 6 *Mordell-Weil rank one rational elliptic surfaces is infinity. In fact, the isomorphism classes depend on a single parameter.*

Lemma 5.5 *The No.3 Mordell-Weil rank one rational elliptic surface S is uniquely determined by its singular fibre type $(I_4, I_2, I_2, I_2, I_2)$, up to isomorphisms.*

Proof. To see the uniqueness of S, we need to calculate the polynomial g_2, g_3 and Δ in the homogeneous coordinates (X, Y) of $\mathbf{P}^1$ where g_2, g_3 are the sections which determine the *Weierstrass model* of S (cf. Theorem 2.11). For simplicity, we let $G_2 = g_2$ and $G_3 = 3\sqrt{3}g_3$. Then $\Delta = G_2^3 - G_3^2$.

The orders ν_p of zeros for the fibre type $(I_4, I_2, I_2, I_2, I_2)$ (cf. [Her]) have to be:

Table 5.2

p	$\nu_p(G_2)$	$\nu_p(G_3)$	$\nu_p(\Delta)$
p_1	0	0	4
p_2	0	0	2
p_3	0	0	2
p_4	0	0	2
p_5	0	0	2
sum	0	0	12

Therefore

$$-\bar{\Delta}^2 = \Delta = G_2^3 - G_3^2 \tag{1}$$

with $\bar{\Delta}$, $G_3 \in H^0(\mathbf{P}^1, \mathcal{O}(6L))$ where $degL = p_g - q + 1 = 1$ since S is a rational surface. Thus (1) is equivalent to

$$G_2^3 = G_3^2 - \bar{\Delta}^2 = (G_3 - \bar{\Delta})(G_3 + \bar{\Delta}).$$

It follows from Table 5.2 that G_2, G_3 and $\bar{\Delta}$ are relatively prime. We let

$$H_1^3 = G_3 - \bar{\Delta} \quad \text{and} \quad H_2^3 = G_3 + \bar{\Delta}$$

It is easy to see that H_1, $H_2 \in H^0(\mathbf{P}^1, \mathcal{O}(2L))$ and they are relatively prime. Thus

$$-\bar{\Delta} = \frac{1}{2}(H_1^3 - H_2^3) = \frac{1}{2}(H_1 - H_2)(\eta^2 H_2 - \eta H_1)(\eta H_2 - \eta^2 H_1)$$

where $\eta = e^{\frac{2\pi i}{3}}$. By (1), $\bar{\Delta}$ has one double zero at p_1 and four single zeros at p_i where i = 2,3,4,5. Let

$$\begin{aligned} J_1 &:= H_1 - H_2 \\ J_2 &:= \eta^2 H_2 - \eta H_1 \\ J_3 &:= \eta H_2 - \eta^2 H_1 \end{aligned}$$

with $J_i \in H^0(\mathbf{P}_1, \mathcal{O}(2L))$, $i = 1, 2, 3$. Similarly, J_1, J_2 and J_3 are relatively prime.

It can be assumed that double point of $\bar{\Delta}$ is that of J_2 at $p_1 = 0$, two of the four single zeros of $\bar{\Delta}$ are that of J_3 at $p_2 = \infty$ and $p_3 = 1$. Therefore we may write

$J_2 = X^2$ and $J_3 = Y(X-Y)$. It follows

$$\begin{aligned} J_1 &= X^2 + XY - Y^2 \\ H_1 &= \frac{1}{1-\eta}(X^2 - \eta XY + \eta Y^2) \\ H_2 &= \frac{1}{1-\eta}(\eta X^2 - XY + Y^2) \\ \bar{\Delta} &= -\frac{1}{2}(Y^2 - XY - X^2)\cdot X^2 \cdot Y(X-Y) \end{aligned}$$

and

$$\begin{aligned} G_2 &= H_1 \cdot H_2 \\ G_3 &= H_1^3 + \bar{\Delta} \\ \Delta &= -\frac{1}{4}(Y^2 - XY - X^2)^2 \cdot X^4 \cdot Y^2(X-Y)^2 \end{aligned}$$

This proves the lemma.

Thus we get the following lemma (cf. the proof of Lemma 3.3):

Lemma 5.6 *Let V be a Picard number two relatively minimal Gorenstein log del Pezzo surface with $K_V^2 = 1$. Then the singularity type of V and the number m of the isomorphism classes of the given singularity type are listed in Table* 5.3.

Table 5.3

Dynkin type of V	m	*Dynkin type of V*	m
D_7	∞	$D_5 + 2A_1$	∞
$A_3 + 4A_1$	1	$D_4 + A_3$	∞

Next we deal with $K_V^2 = 2$.

By blowing down an NEC on U_1, we get a surface U_2 which is the minimal resolution of the Picard number two Gorenstein log del Pezzo surface V with $K_V^2 = 2$.

Theorem 5.7 *Let V be the Picard number two Gorenstein log del Pezzo surface with $K_V^2 = 2$.*
(1) *There exist* 12 *singularity types which are listed in the following: E_6, $D_4 + 2A_1$, $A_4 + A_2$, $A_3 + 3A_1$, D_6, A_6, $2A_3$, $3A_2$, $D_5 + A_1$, $A_5 + A_1$, $A_3 + A_2 + A_1$, $6A_1$.*
(2) *If V is a relatively minimal surface, then the singularity type of V and the number m of the isomorphism classes with the given singularity type are listed in Table* 5.4.

Table 5.4

Dynkin type of V	m	*Dynkin type of V*	m
D_6	∞	$D_4 + 2A_1$	1
$6A_1$	1	$2A_3$	∞

Proof. The singularity types and the relatively minimal singularity types can be found by a direct calculation together with the definition of the relatively minimal surfaces (cf. Definition 2.1). For the details, we refer to Table 3 of Appendix.

Now we shall deal with the number of isomorphism classes with the given relatively minimal singularity type separately. Firstly, we prove the following lemma.

Lemma 5.8 *The number of the isomorphism classes of the Picard number two Gorenstein log del Pezzo surface with the singularity type D_6 is infinity.*

Proof. We let U be the minimal resolution of V. By Table 3 of Appendix, there are four possibilities for U: start from No.1 (2, 7 or 8) Mordell-Weil rank one rational elliptic surface. Since each of these surfaces has at least one type I_1 fibre, we have

Claim 5.9 *There exists a reduced irreducible nodal member $A \in |-K_U|$.*

From the configuration of the negative curves in U which is listed below, we know there are three (-1)-curves: l_1, l_2 and l_3.



Figure 5.1

Claim 5.10 *There exists an automorphism on U which interchanges l_1 and l_2.*

Proof of the claim. Let $W \longrightarrow U$ be the blow up of $p = A \cap l_3$. It is easy to see that W is a surface which is the minimal resolution of the Picard number two Gorenstein log del Pezzo surface with the singularity type E_7. Thus W is either $U_1^7(E_7)$ or $U_1^8(E_7)$ and the proper transform of l_i (i=1,2) intersects the 1st component of E_7. From Table 2 of Appendix, we know that l_i (i=1,2) comes from the Mordell-Weil section P_1 or P_{-1} of the No.7 or 8 Mordell-Weil rank one rational elliptic surface where P_1 is the generator of the rank one Mordell-Weil lattice A_1^*. In any case, we may define an automorphism Φ on No.7 (No.8) Mordell-Weil rank one rational elliptic surface which is the involution on a general fibre. Then $\Phi(P_1) = P_{-1}$ and Φ will induce an automorphism on U which interchanges l_1 and l_2. This proves the claim.

Let $U \longrightarrow \mathbf{P}^2$ be the composition of blowing down of $l_3 + G_2 + G_6 + G_3$, $l_1 + G_1 + G_4$ to p, q respectively where G_i $(i = 1, \cdots, 6)$ is the component of D_6 (cf. Figure 5.1).



Figure 5.2

From the above construction and Claim 5.10, we know that U is uniquely determined by a pair $M = (S, C, p)$ where $S = \mathbf{P}^2$, C is a nodal cubic curve on S and p is a point of C which is neither a singular point of C nor an inflex of C. An automorphism of U will induce an isomorphism on $M = (S, C, p)$ by making use of Claim 5.10 if necessary. In order to prove the lemma, we have only to show that there exist infinitly many isomorphism classes of such pairs.

Before stating the next result, we employ the following notation.
$S = \mathbf{P}^2$ with coordinates X,Y and Z.
$C = \{[X : Y : Z] | Y^2 Z = X^2(X + Z)\}$.
$S' = \mathbf{P}^2$ with coordinates X',Y' and Z'.
$C' = \{[X' : Y' : Z'] | Y'^2 Z' = X'^2(X' + Z')\}$.

Claim 5.11 *If* $\Phi : S \longrightarrow S'$ *is an isomorphism such that* $\Phi(C) = C'$*, then either* $\Phi([X : Y : Z]) = [X' : Y' : Z']$ *or* $\Phi([X : Y : Z]) = [X' : -Y' : Z']$.

Proof of the claim. Clearly, Φ is linear map. We set:

$$\begin{pmatrix} X' \\ Y' \\ Z' \end{pmatrix} = \Phi \begin{pmatrix} X \\ Y \\ Z \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \cdot \begin{pmatrix} X \\ Y \\ Z \end{pmatrix} = A \cdot \begin{pmatrix} X \\ Y \\ Z \end{pmatrix}$$

Then $det A \neq 0$. Since $\Phi(C) = C'$, we have

$$\Phi([0:0:1]) = [0:0:1] \quad \text{and} \quad \Phi([1:0:-1]) = [1:0:-1].$$

From the above equalities, we may set

$$a_{33} = 1, a_{11} = 1 - a_{31}, a_{13} = a_{23} = a_{21} = 0.$$

For simplicity, we denote $a_{12} = a$, $a_{22} = b$, $a_{32} = c$ and $a_{11} = d$. Thus we have

$$X' = dX + aY, \quad Y' = bY, \quad Z' = (1 - d)X + cY + Z.$$

Substituting above equalities to $Y'^2 Z' = X'^2(X' + Z')$, we get

$$(bY)^2 \cdot ((1 - d)X + cY + Z) = (dX + aY)^2 \cdot (X + aY + cY + Z). \tag{2}$$

(2) is hold for all $[X : Y : Z] \in C$. On the other hand, if $[X : Y : Z] \in C$, then $[X : -Y : Z] \in C$, thus we also have

$$(bY)^2 \cdot ((1-d)X - cY + Z) = (dX - aY)^2 \cdot (X - aY - cY + Z). \tag{3}$$

From (2) + (3), we may get

$$\begin{aligned} b^2Y^2[(1-d)X + Z] &= (X+Z)(d^2X^2 + a^2Y^2) + 2ad(a+c)Y^2X \\ &= d^2Y^2Z + a^2Y^2(X+Z) + 2ad(a+c)XY^2. \end{aligned}$$

So for all $[X : Y : Z] \in C - \{[0 : 0 : 1], [1 : 0 : -1]\}$, we must have

$$b^2[(1-d)X + Z] = d^2Z + a^2(X+Z) + 2ad(a+c)X$$

Since we can take $Z = 1$ and arbitrarily many X, we get

$$b^2 = d^2 + a^2 \tag{4}$$

$$b^2(1-d) = a^2 + 2ad(a+c) \tag{5}$$

By (2) − (3), we get

$$cb^2Y^3 = 2adXY(X+Z) + (a+c)d^2YX^2 + a^2(a+c)Y^3.$$

Since above equality holds for all $[X : Y : Z] \in C - \{[0 : 0 : 1], [1 : 0 : -1]\}$, we get

$$cb^2Y^2 = 2adX(X+Z) + (a+c)d^2X^2 + a^2(a+c)Y^2.$$

We let $g(X,Y,Z) = cb^2Y^2 - [2adX(X+Z) + (a+c)d^2X^2 + a^2(a+c)Y^2]$. Then we know C is included in the closure of the zero set of g. On the other hand, C is an irreducible cubic. Thus we know $g = 0$ and so

$$2ad = 0, \tag{6}$$

$$b^2c = a^2(a+c). \tag{7}$$

By (6), we have $a = 0$ or $d = 0$. But $detA \neq 0$, we see that a and d can't both equal to 0. If $d = 0$, then by (4) and (7), we get $a = 0$, a contradiction. Thus $a = 0$. Then by (5) and $detA \neq 0$ which implies $b \neq 0$, we have $d = 1$ and so by (4), $b = \pm 1$. This proves the claim.

Let $M = (S, C, p_1)$ and $M_1 = (S, C, p_2)$ where S and C are defined above and $p_1 = [X_1 : Y_1 : Z_1]$ and $p_2 = [X_2 : Y_2 : Z_2]$. From the above claim, we see that M_1 and M_2 are not isomorphic if $[X_1 : Y_1 : Z_1] \neq [X_2 : \pm Y_2 : Z_2]$. Moving $p \in C$, we may get infinitely many non-isomorphism classes of the pair (S, C, p) and so the infinite many non-ismorphism classes of U. This proves the lemma.

Next we consider the singularity type $6A_1$.

From Table 3 of the Appedenix, there are at most two possibilities for $U(D_6)$: Start with either $U_1^3(A_3+4A_1)$ or $U_1^{17}(D_4+3A_1)$ and blow down an NEC. Since the configuration of $U_2^3(6A_1)$ and $U_2^{17}(6A_1)$ are the same, we only discuss one of these, say, $U_2^3(6A_1)$. By Tables 1 and 2 in the Appedenix, we find that blowing up a point on Q_1 or Q_2 will give rise to the surface U_1 which is the minimal resolution of a Picard number two log del Pezzo surface with singularity type A_3+4A_1. By Lemma 5.6, we know $U_1 = U_1^3(A_3+4A_1)$. On the other hand, there is a unique way to blow down an NEC on $U_1^3(A_3+4A_1)$ to $U_2^3(6A_1)$, modular an automorphism switching the two NECs (as in Claim 5.10). Thus the uniqueness of the surface with the singularity type $6A_1$ is due to the uniqueness of the Picard number two log del Pezzo surface with the singularity type A_3+4A_1.

Now we deal with the singularity type D_4+2A_1.

We let $V(D_4+2A_1)$ be the Picard number two Gorenstein log del Pezzo surface with the singularity type D_4+2A_1, $U(D_4+2A_1)$ the minimal resolution of $V(D_4+2A_1)$ and D the exceptional divisor. The configuration of D is in the left part of the following figure. Blowing up p, q of D will give rise to the No.9 extremal rational elliptic surface Y with the singular fibre type (I_1^*, I_4, I_1) which is unique up to isomorphisms (cf. [MP, Theorem 5.4]). On the other hand, a direct calculation shows that the Mordell-Weil group of Y is $\mathbf{Z}/4\mathbf{Z}$ with the generator Q and $Y \longrightarrow U(D_4+2A_1)$ is the blow-down of $\mathcal{O}$ and $2Q$ on Y.



Figure 5.3

Finally, we look at the singularity type $2A_3$.

We let $V(2A_3)$ be the Picard number two Gorenstein log del Pezzo surface with the singularity type $2A_3$, $U(2A_3)$ the minimal resolution of $V(2A_3)$ and D the exceptional divisor. The configuration of D is in the left part of the following figure.



Figure 5.4

We denote

A = the set of the isomorphism classes of $U(2A_3)$.
B = the set of the extremal rational elliptic surfaces with the singular fibre type (I_0^*, I_0^*).

We define an surjective map $\Phi : A \longrightarrow B$ where Φ is the blow-up p and q of D. Thus the infinity of the number of the isomorphism classes of $U(2A_3)$ is due to the infinity of the number of the isomorphism classes of the extremal rational elliptic surface with the singular fibre type (I_0^*, I_0^*) (cf. [MP, Theorem 5.4]).

We may do the same operation as above to get a surface which is the minimal resolution of the Picard number two Gorenstein log del Pezzo surface V with $K_V^2 \geq 3$. For details, we refer to the Table 3 of the Appedenix.

Lemma 5.12 *Let V be the Picard number two Gorenstein log del Pezzo surface with $K_V^2 \geq 3$. Then the singularity type of V is one of the following listed in Table* 5.5.

Table 5.5

K_V^2	*Dynkin type of V*
3	D_5, A_5, $A_4 + A_1$, $A_3 + 2A_1$, $2A_2 + A_1$
4	D_4, A_4, $A_3 + A_1$, $A_2 + 2A_1$, $4A_1$
5	A_3, $A_2 + A_1$
6	A_2, $2A_1$
7	A_1

Remark 5.13 *Let V be a Picard number two Gorenstein log del Pezzo surface and U the minimal resolution of V.*

(1) *From a direct calulation, we find that the configurations of the negative curves with the same singularity type are the same except for the type $A_5 + A_1$ which has two different configurations where one has two NECs (cf. $U_2^9(A_5 + A_1)$) and the other has three NECs (cf. $U_2^{10}(A_5 + A_1)$ or $U_2^{11}(A_5 + A_1)$).*

(2) *The configurations of the negative curves of U with $K_U^2 \geq 3$ are listed in the Figure* 6 *of Appendix. From these configurations, it is easy to see that there are only two relatively minimal Dynkin types in the above table which are $4A_1$ and D_4.*

Lemma 5.14 *The Picard number two relatively minimal Gorenstein log del Pezzo surface with the singularity type* $4A_1$ *is unique.*

Proof. We let U be the minimal resolution of $V(4A_1)$ and D the exceptional divisor. Then the configuation of D is in the left part of the following figure.



Figure 5.5

Let $\sigma_1 : U \longrightarrow W$ be the contraction of l_1 and l_2 and $\sigma_2 : W \longrightarrow \mathbf{P}^1 \times \mathbf{P}^1$ the contraction of E_1 and E_2. By changing the coordinates of $\mathbf{P}^1 \times \mathbf{P}^1$, we may assume $\sigma_2 \circ \sigma_1(E_1) = p_1 = (\infty, \infty)$ and $\sigma_2 \circ \sigma_1(E_2) = p_2 = (0, 0)$. There is unique way from $(\mathbf{P}^1 \times \mathbf{P}^1, p_1, p_2)$ to U. Thus we prove the lemma.

Lemma 5.15 *The Picard number two relatively minimal Gorenstein log del Pezzo surface with the singularity type* D_4 *is unique.*

Proof. From Figure 6 in Appendix, we know that there is unique way from the surface $U(D_5)$ which is the minimal resolution of the Picard number two Gorenstein log del Pezzo surface with the singularity type D_5 to $U(D_4)$.

Let $U(A_1 + D_6) \longrightarrow U(D_5)$ be the blow up at the point p (see Figure 6 of Appendix). Then $U(A_1 + D_6)$ is the minimal resolution of a Picard number one Gorenstein log del Pezzo surface with the singularity type $A_1 + D_6$. On the other hand, $U(A_1 + D_6)$ is unique by Lemma 4.6 and there is a unique way from $U(A_1 + D_6)$ to $U(D_5)$. Thus we know $U(D_5)$ is unique up to isomorphisms. This proves the lemma.

Combining Lemma 5.1, Theorem 5.7 and Lemma 5.12, we prove Theorem 1.5.

Combining Lemmas 5.6, Theorem 5.7, Lemmas 5.14 and 5.15, we prove Theorem 1.6.

Appendix. List of Configurations and Tables

Configurations which is used in the proof of Theorem 1.2

In the following configurations, we exclude the fibres of type I_1 or II which does not contribute to the calculation of the Mordell-Weil group of the corresponding fibration

[Sh]. For simplicity, we only list the (-1)-curves which will be used in the proof of Theorem 1.2.

——————— Stands for a (-2)-curve

- - - - - - - - - - - - Stands for a (-1)-curve

·························· Stands for a curve with self-intersection 0.



Figure 1





Figure 2



Figure 3



Figure 4

Configurations of the Singularity types

(a) The Picard number one Gorenstein log del Pezzo surfaces.

The following configurations are the figures of the negative curves on U which is the minimal resolution of the Picard number one Gorenstein log del Pezzo surface and $2 \leq K_U^2 \leq 7$.



E_7 $A_1 + D_6$ A_7

$A_2 + A_5$ $3A_1 + D_4$ $A_1 + 2A_3$

$3A_2$ $A_1 + A_5$ E_6

$2A_1 + A_3$ D_5 A_4 $A_2 + A_1$

Figure 5

(b) The Picard number two Gorenstein log del Pezzo surface.

The following configurations are the figures of the negative curves on U which is the minimal resolution of the Picard number two Gorenstein log del Pezzo surface and $K_U^2 \geq 3$.



D_5



A_5



$A_4 + A_1$



A_3+2A_1 $2A_2+A_1$ D_4

A_4 A_3+A_1 A_2+2A_1 $4A_1$

A_3 A_2+A_1 A_2

$2A_1$ A_1

Figure 6

Table 1. Mordell-Weil Rank one Rational Elliptic Surfaces

In the following table, $Y \longrightarrow U_1(D)$ is the blow-down of the zero section $\mathcal{O}$ and $U_1(D) \longrightarrow V$ is the minimal resolution of the singularities of V with D the exceptional divisor; this map is nothing but the contraction of all (-2)-curves. The symble $\langle m\rangle$ stands for a rank one lattice $\mathbf{Z}P$ with the generator P such that $\langle P,P\rangle = m$. We also denote $P_n := nP$ where $n \in \mathbf{Z}$. The A_1^* indicates the lattice generated by the element P such that $< P,P >= 1/2$. In the last column, E is a (-1)-curve and R is a connected component of D such that $E+R$ is a linear chain on $U_1(D)$. We can't find such a linear chain for the first 6 cases.

♯	Fibre type on Y	MW(Y)	$U_1(D)$	$E+R$ on $U_1(D)$
1	I_3^*, II, I_1	$< 1/4 >$	$U_1^1(D_7)$	
2	$I_3^*, 3I_1$	$< 1/4 >$	$U_1^2(D_7)$	
3	$I_4, 4I_2$	$< 1/4 > \oplus(\mathbf{Z}/2\mathbf{Z})^2$	$U_1^3(A_3+4A_1)$	
4	I_1^*, III, I_2	$< 1/4 > \oplus(\mathbf{Z}/2\mathbf{Z})$	$U_1^4(D_5+2A_1)$	
5	$I_1^*, 2I_2, I_1$	$< 1/4 > \oplus(\mathbf{Z}/2\mathbf{Z})$	$U_1^5(D_5+2A_1)$	
6	$I_0^*, I_4, 2I_1$	$< 1/4 > \oplus(\mathbf{Z}/2\mathbf{Z})$	$U_1^6(D_4+A_3)$	
7	III^*, II, I_1	A_1^*	$U_1^7(E_7)$	P_2
8	$III^*, 3I_1$	A_1^*	$U_1^8(E_7)$	P_2
9	$I_8, 4I_1$	$A_1^* \oplus (\mathbf{Z}/2\mathbf{Z})$	$U_1^9(A_7)$	P_2
10	$I_8, II, 2I_1$	$< 1/8 >$	$U_1^{10}(A_7)$	P_3+A_7
11	$I_8, 4I_1$	$< 1/8 >$	$U_1^{11}(A_7)$	P_3+A_7
12	$2I_4, I_2, 2I_1$	$A_1^* \oplus (\mathbf{Z}/4\mathbf{Z})$	$U_1^{12}(2A_3+A_1)$	P_2
13	I_2^*, III, I_1	$A_1^* \oplus \mathbf{Z}/2\mathbf{Z}$	$U_1^{13}(D_6+A_1)$	P_2
14	$I_2^*, I_2, 2I_1$	$A_1^* \oplus \mathbf{Z}/2\mathbf{Z}$	$U_1^{14}(D_6+A_1)$	P_2
15	$I_6, IV, 2I_1$	$A_1^* \oplus \mathbf{Z}/3\mathbf{Z}$	$U_1^{15}(A_5+A_2)$	P_2
16	$I_6, I_3, 3I_1$	$A_1^* \oplus \mathbf{Z}/3\mathbf{Z}$	$U_1^{16}(A_5+A_2)$	P_2
17	$I_0^*, 3I_2$	$A_1^* \oplus (\mathbf{Z}/2\mathbf{Z})^2$	$U_1^{17}(D_4+3A_1)$	P_2
18	IV^*, III, I_1	$< 1/6 >$	$U_1^{18}(E_6+A_1)$	P_3+A_1
19	IV^*, I_2, II	$< 1/6 >$	$U_1^{19}(E_6+A_1)$	P_3+A_1
20	$IV^*, I_2, 2I_1$	$< 1/6 >$	$U_1^{20}(E_6+A_1)$	P_3+A_1
21	I_4, I_3, III, I_2	$< 1/12 > \oplus\mathbf{Z}/2\mathbf{Z}$	$U_1^{21}(A_3+A_2+2A_1)$	P_4+A_2
22	$I_4, I_3, 2I_2, I_1$	$< 1/12 > \oplus\mathbf{Z}/2\mathbf{Z}$	$U_1^{22}(A_3+A_2+2A_1)$	P_4+A_2
23	$IV, 2I_3, I_2$	$< 1/6 > \oplus\mathbf{Z}/3\mathbf{Z}$	$U_1^{23}(3A_2+A_1)$	P_3+A_1
24	$3I_3, I_2, I_1$	$< 1/6 > \oplus\mathbf{Z}/3\mathbf{Z}$	$U_1^{24}(3A_2+A_1)$	P_3+A_1
25	$I_7, III, 2I_1$	$< 1/14 >$	$U_1^{25}(A_6+A_1)$	P_4+A_6
26	I_7, I_2, II, I_1	$< 1/14 >$	$U_1^{26}(A_6+A_1)$	P_4+A_6
27	$I_7, I_2, 3I_1$	$< 1/14 >$	$U_1^{27}(A_6+A_1)$	P_4+A_6
28	I_1^*, IV, I_1	$< 1/12 >$	$U_1^{28}(D_5+A_2)$	P_4+A_2
29	I_1^*, I_3, II	$< 1/12 >$	$U_1^{29}(D_5+A_2)$	P_4+A_2
30	$I_1^*, I_3, 2I_1$	$< 1/12 >$	$U_1^{30}(D_5+A_2)$	P_4+A_2
31	I_6, III, I_2, I_1	$< 1/6 > \oplus\mathbf{Z}/2\mathbf{Z}$	$U_1^{31}(A_5+2A_1)$	P_3+A_1
32	$I_6, 2I_2, 2I_1$	$< 1/6 > \oplus\mathbf{Z}/2\mathbf{Z}$	$U_1^{32}(A_5+2A_1)$	P_3+A_1
33	I_5, I_4, II, I_1	$< 1/20 >$	$U_1^{33}(A_4+A_3)$	P_5+A_3
34	$I_5, I_4, 3I_1$	$< 1/20 >$	$U_1^{34}(A_4+A_3)$	P_6+A_4
35	I_5, IV, I_2, I_1	$< 1/30 >$	$U_1^{35}(A_4+A_2+A_1)$	P_6+A_4
36	I_5, I_3, III, I_1	$< 1/30 >$	$U_1^{36}(A_4+A_2+A_1)$	P_6+A_4
37	I_5, I_3, I_2, II	$< 1/30 >$	$U_1^{37}(A_4+A_2+A_1)$	P_6+A_4
38	$I_5, I_3, I_2, 2I_1$	$< 1/30 >$	$U_1^{38}(A_4+A_2+A_1)$	P_6+A_4

Table 2. Nice Exceptinal Curves on U_1

In the following table, we employ the following notation and convention:

We let G_i, H_i, J_i, L_i and M_i be the i-th component in the corresponding fibres F_1, F_2, F_3 F_4 and F_5 respectively. The numbering of the singular fibre is defined in the following diagrams. For the three types of singular fibres, the 0-th component in a fibre intersects the zero section. For the other types, the numbering is natural. The components of the singular fibre are numbered as in [Ye]. For a lattice $L =< m > \oplus \mathbf{Z}/k\mathbf{Z}$, we let P_1 to be the generator of $< m >$ and $P_n = nP_1$ where $n \in \mathbf{Z}$. If $Q \in \mathbf{Z}/k\mathbf{Z}$, i.e., the torsion part of L, then we denote $P_{n,Q} = P_n + Q$. (R_1R_2) is the intersection number of the sections R_1 and R_2.

$\sharp$	Fibre type on Y	Sections disjoint from $\mathcal{O}$ on Y	$NECs$ on U_1	sections which intersects NEC on U_1
1	I_3^*, II, I_1	$P_1 = (2)$, $P_{-1} = (3)$	$P_{\pm 1}$, $P_{\pm 2}$	$(P_{\pm 1}P_{\mp 2}) = 1$
2	$I_3^*, 3I_1$	$P_{\pm 2} = (1)$	G_0	$(P_2P_{-2}) = 1$
3	$I_4, 4I_2$	$Q_1 = (2,1,1,0,0)$, $Q_2 = (2,0,0,1,1)$, $Q_3 = (0,1,1,1,1)$, $P_1 = (1,0,1,1,0)$, $P_{1,Q_1} = (3,1,0,1,0)$, $P_{1,Q_2} = (3,0,1,0,1)$, $P_{1,Q_3} = (1,1,0,0,1)$, $P_{-1} = (1,0,1,0,1)$, $P_{-1,Q_1} = (3,1,0,0,1)$, $P_{-1,Q_2} = (3,0,1,1,0)$, $P_{-1,Q_3} = (1,1,0,1,0)$, $P_{\pm 2} = (2,0,0,0,0)$, $P_{\pm 2,Q_1} = (0,1,1,0,0)$, $P_{\pm 2,Q_2} = (0,0,0,1,1)$.	$P_{\pm 2}$	$(P_2P_{-2}) = 1$ $(P_{\pm 2}P_{\mp 2,Q_1}) = 2$ $(P_{\pm 2}P_{\mp 1}) = 1$ $(P_{\pm 2}P_{\mp 1,Q_2}) = 1$ $(P_{\pm 2}P_{\mp 1,Q_1}) = 1$ $(P_{\pm 2}P_{\mp 1,Q_2}) = 1$ $(P_{\pm 2}Q_3) = 1$

4 5	I_1^*, III, I_2 $I_1^*, 2I_2, I_1$	$Q = (1,1,1),\ P_1 = (2,1,0),$ $P_{1,Q} = (3,0,1),\ P_{-1} = (2,0,1),$ $P_{-1,Q} = (3,1,0),\ P_{\pm 2} = (1,0,0),$ $P_{\pm 2,Q} = (0,1,1).$	$P_{\pm 2}$ G_0	$(P_{\pm 2} P_{\mp 1}) = 1$ $(P_{\pm 2} P_{\mp 1,Q}) = 1$ $(P_{\pm 2} P_{\mp 2,Q}) = 2$ $(P_2 P_{-2}) = 1$ $(G_0 P_{\pm 2,Q}) = 1$
6	$I_0^*, I_4, 2I_1$	$Q = (1,2),\ P_1 = (2,1)$ $P_{1,Q} = (3,3),\ P_{-1} = (3,1)$ $P_{-1,Q} = (2,3),\ P_{\pm 2} = (1,0)$ $P_{\pm 2,Q} = (0,2)$	$P_{\pm 2},$ $P_{\pm 2,Q}$ G_0	$(P_{\pm 2} P_{\mp 1}) = 1$ $(P_{\pm 2} P_{\mp 1,Q}) = 1$ $(P_2 P_{-2}) = 1$ $(P_{\pm 2,Q} P_{\mp 1}) = 1$ $(P_{\pm 2} P_{\mp 2,Q}) = 2$ $(P_{2,Q} P_{-2,Q}) = 1$ $(G_0 P_{\pm 2,Q}) = 1$
7 8	III^*, II, I_1 $III^*, 3I_1$	$P_{\pm 1} = (1),\ P_{\mp 2} = (0)$	$P_{\pm 1}$ G_0	
9	$I_8, 4I_1$	$Q = (4),\ P_{\pm 1} = (2)$ $P_{\pm 1,Q} = (6),\ P_{\pm 2} = (0)$	$Q,\ P_{\pm 1}$ $P_{\pm 1,Q}$	$(P_{\pm 1} P_{\mp 1,Q}) = 1$ $(P_{\pm 2} Q) = 1$
10 11	$I_8, II, 2I_1$ $I_8, 4I_1$	$P_1 = (3),\ P_2 = (6)$ $P_3 = (1),\ P_{-1} = (5)$ $P_{-2} = (2),\ P_{-3} = (7)$	$P_{\pm 1}$ $P_{\pm 2}$ $P_{\pm 3}$	$(P_{\pm 1} P_{\mp 3}) = 1$ $(P_2 P_{-2}) = 1$ $(P_{\pm 3} P_{\mp 2}) = 1$
12	$2I_4, I_2, 2I_1$	$Q_1 = (1,1,1),\ Q_2 = (2,2,0)$ $Q_3 = (3,3,1),\ P_{\pm 1} = (2,0,1)$ $P_{\pm 1,Q_1} = (3,1,0),\ P_{\pm 1,Q_2} = (0,2,1)$ $P_{\pm 1,Q_3} = (1,3,0),\ P_{\pm 2} = (0,0,0)$	$\sharp\{NEC\}$ $= \phi$	
13 14	I_2^*, III, I_1 $I_2^*, I_2, 2I_1$	$Q = (2,1),\ P_{\pm 1} = (3,0)$ $P_{\pm 1,Q} = (1,1),\ P_{\pm 2} = (0,0)$	$P_{\pm 1}$ G_0	$(P_{\pm 1} P_{\mp 1,Q}) = 1$
15 16	$I_6, IV, 2I_1$ $I_6, I_3, 3I_1$	$Q_1 = (2,1),\ Q_2 = (4,2)$ $P_{\pm 1} = (3,0),\ P_{\pm 1,Q_1} = (5,1)$ $P_{\pm 1,Q_2} = (1,2),\ P_{\pm 2} = (0,0)$	$P_{\pm 1}$	$(P_{\pm 1} P_{\mp 1,Q_1}) = 1$ $(P_{\pm 1} P_{\mp 1,Q_2}) = 1$
17	$I_0^*, 3I_2$	$Q_1 = (1,1,1,0),\ Q_2 = (2,0,1,1)$ $Q_3 = (3,1,0,1),\ P_{\pm 1} = (0,1,1,1)$ $P_{\pm 1,Q_1} = (1,0,0,1),\ P_{\pm 1,Q_2} = (2,1,0,0)$ $P_{\pm 1,Q_3} = (3,0,1,0),\ P_{\pm 2} = (0,0,0,0)$	G_0	$(G_0 P_{\pm 1}) = 1$ $(G_0 P_{\pm 2}) = 1$
18 19 20	IV^*, III, I_1 IV^*, I_2, II $IV^*, I_2, 2I_1$	$P_1 = (1,1),\ P_2 = (2,0)$ $P_{\pm 3} = (0,1),\ P_{-1} = (2,1)$ $P_{-2} = (1,0)$	$P_{\pm 2}$ $P_{\pm 3}$ G_0	$(P_{\pm 3} P_{\mp 1}) = 1$ $(P_2 P_{-2}) = 1$ $(P_{\pm 3} P_{\mp 2}) = 2$ $(P_3 P_{-3}) = 2$ $(G_0 P_{\pm 3}) = 1$
21 22	I_4, I_3, III, I_2 $I_4, I_3, 2I_2, I_1$	$Q = (2,0,1,1),\ P_1 = (1,1,1,0)$ $P_{1,Q} = (3,1,0,1),\ P_2 = (2,2,0,0)$ $P_{2,Q} = (0,2,1,1),\ P_3 = (3,0,1,0)$ $P_{3,Q} = (1,0,0,1),\ P_4 = (0,1,0,0)$ $P_{-1} = (1,2,0,1),\ P_{-1,Q} = (3,2,1,0)$ $P_{-2} = (2,1,0,0),\ P_{-2,Q} = (0,1,1,1)$ $P_{-3} = (3,0,0,1),\ P_{-3,Q} = (1,0,1,0)$ $P_{-4} = (0,2,0,0)$	$P_{\pm 4}$	$(P_{\pm 4} P_{\mp 1}) = 1$ $(P_{\pm 4} P_{\mp 1,Q}) = 1$ $(P_{\pm 4} P_{\mp 2}) = 1$ $(P_{\pm 4} P_{\mp 2,Q}) = 1$ $(P_{\pm 4} P_{\mp 3}) = 2$ $(P_{\pm 4} P_{\mp 3,Q}) = 2$ $(P_4 P_{-4}) = 2$

23 24	$IV, 2I_3, I_2$ $3I_3, I_2, I_1$	$Q_1 = (1,1,1,0)$, $Q_2 = (2,2,2,0)$ $P_1 = (1,2,0,1)$, $P_{1,Q_1} = (2,0,1,1)$ $P_{1,Q_2} = (0,1,2,1)$, $P_2 = (1,0,2,0)$ $P_{2,Q_1} = (2,1,0,0)$, $P_{2,Q_2} = (0,2,1,0)$ $P_{-1} = (1,0,2,1)$, $P_{-1,Q_1} = (2,1,0,1)$ $P_{-1,Q_2} = (0,2,1,1)$, $P_{-2} = (1,2,0,0)$ $P_{-2,Q_1} = (2,0,1,0)$, $P_{-2,Q_2} = (0,1,2,0)$ $P_{\pm 3} = (0,0,0,1)$	$P_{\pm 3}$	$(P_{\pm 3}P_{\mp 1}) = 1$ $(P_{\pm 3}P_{\mp 1,Q_1}) = 1$ $(P_{\pm 3}P_{\mp 1,Q_2}) = 1$ $(P_{\pm 3}P_{\mp 2}) = 2$ $(P_{\pm 3}P_{\mp 2,Q_1}) = 2$ $(P_{\pm 3}P_{\mp 2,Q_2}) = 2$ $(P_3P_{-3}) = 2$
25 26 27	$I_7, III, 2I_1$ I_7, I_2, II, I_1 $I_7, I_2, 3I_1$	$P_1 = (2,1)$, $P_2 = (4,0)$ $P_3 = (6,1)$, $P_4 = (1,0)$ $P_{-1} = (5,1)$, $P_{-2} = (3,0)$ $P_{-3} = (1,1)$, $P_{-4} = (6,0)$	$P_{\pm 2}$ $P_{\pm 4}$	$(P_{\pm 3}P_{\mp 2}) = 1$ $(P_{\pm 4}P_{\mp 1}) = 1$ $(P_{\pm 4}P_{\mp 2}) = 1$ $(P_{\pm 4}P_{\mp 3}) = 1$ $(P_4P_{-4}) = 2$
28 29 30	I_1^*, IV, I_1 I_1^*, I_3, II $I_1^*, I_3, 2I_1$	$P_1 = (2,1)$, $P_2 = (1,2)$ $P_3 = (3,0)$, $P_4 = (0,1)$ $P_{-1} = (3,2)$, $P_{-2} = (1,1)$ $P_{-3} = (2,0)$, $P_{-4} = (0,2)$	$P_{\pm 3}$ $P_{\pm 4}$ G_0	$(P_{\pm 3}P_{\mp 2}) = 1$ $(P_{\pm 4}P_{\mp 2}) = 1$ $(P_{\pm 4}P_{\mp 3}) = 2$ $(P_{-4}P_4) = 2$ $(P_{-3}P_3) = 1$ $(G_0P_{\pm 4}) = 1$
31 32	I_6, III, I_2, I_1 $I_6, 2I_2, 2I_1$	$Q = (3,1,0)$, $P_1 = (2,0,1)$ $P_{1,Q} = (5,1,1)$, $P_2 = (4,0,0)$ $P_{2,Q} = (1,1,0)$, $P_{\pm 3} = (0,0,1)$ $P_{-1} = (4,0,1)$, $P_{-1} + Q = (1,1,1)$ $P_{-2} = (2,0,0)$, $P_{-2} + Q = (5,1,0)$	$P_{\pm 2}$ $P_{\pm 3}$	$(P_{\pm 3}P_{\mp 1}) = 1$ $(P_2P_{-2}) = 1$ $(P_{\pm 2}P_{\mp 2,Q}) = 1$ $(P_{\pm 3}P_{\mp 2}) = 2$ $(P_{\pm 3}P_{\mp 2,Q}) = 2$ $(P_3P_{-3}) = 2$
33 34	I_5, I_4, II, I_1 $I_5, I_4, 3I_1$	$P_1 = (2,1)$, $P_2 = (4,2)$ $P_3 = (1,3)$, $P_4 = (3,0)$ $P_5 = (0,1)$, $P_{-1} = (3,3)$ $P_{-2} = (1,2)$, $P_{-3} = (4,1)$ $P_{-4} = (2,0)$, $P_{-5} = (0,3)$	$P_{\pm 4}$ $P_{\pm 5}$	$(P_{\pm 4}P_{\mp 3}) = 1$ $(P_{\pm 5}P_{\mp 3}) = 1$ $(P_4P_{-4}) = 1$ $(P_{\pm 5}P_{\mp 4}) = 2$ $(P_5P_{-5}) = 2$
35 36 37 38	I_5, IV, I_2, I_1 I_5, I_3, III, I_1 I_5, I_3, I_2, II $I_5, I_3, I_2, 2I_1$	$P_1 = (1,1,1)$, $P_2 = (2,2,0)$ $P_3 = (3,0,1)$, $P_4 = (4,1,0)$ $P_5 = (0,2,1)$, $P_6 = (1,0,0)$ $P_{-1} = (4,2,1)$, $P_{-2} = (3,1,0)$ $P_{-3} = (2,0,1)$, $P_{-4} = (1,2,0)$ $P_{-5} = (0,1,1)$, $P_{-6} = (4,0,0)$	$P_{\pm 6}$	$(P_{\pm 6}P_{\mp 1}) = 1$ $(P_{\pm 6}P_{\mp 2}) = 1$ $(P_{\pm 6}P_{\mp 3}) = 1$ $(P_{\pm 6}P_{\mp 4}) = 1$ $(P_{\pm 6}P_{\mp 5}) = 2$ $(P_6P_{-6}) = 2$

Table 3. The Singularity types of the Picard number two Gorenstein log del Pezzo surfaces

To find out the singularity types of the Picard number two Gorenstein log del Pezzo surfaces, we start blowing down the zero-section $\mathcal{O}$ of the Mordell-Weil rank one rational elliptic surface (cf. Theorem 3.5) and then blow down an NEC (if exists) on the resulting surface, and continue this process. The following table indicates the all singularity types found by this method.

In this table, the numbers i in the first column indicates that we start from the No.i Mordell-Weil rank one rational elliptic surface (cf. Table 1 of Appendix). Since we pay attention to the singularity types, if there are more than one surfaces with the same Mordell-Weil group which may produce the same singularity type, then we only pick up one of those surfaces. For example, blowing down the zero section of No.1 and No.2 Mordell-Weil rank one rational elliptic surfaces will give rise to the same singularity type D_7 and the same configuration of the negative curves on the surface; thus we may only choose one surface, i.e., the No.1 Mordell-Weil rank one rational elliptic surface. In the last column of the table, the number m in " Exceptional divisor of $U_1 \longrightarrow U_m$" is the largest number m such that U_m is the minimal resolution of the singularities of the Picard number two Gorenstein log del Pezzo surface. For the notation of P_i, G_j, H_k and Q, we refer to the Table 2 of Appendix. The suffix n of U_n singifies that $K_{U_n}^2 = n$.

Let's give an example, say, the type D_7, to explain how to read this table. There are five NECs on $U_1(D_7)$: $P_{\pm 1}$, $P_{\pm 2}$ and G_0 (cf. Table 2 of Appendix). Thus there are five possibilities to do operation (blowing down) on $U_1(D_7)$. But by the symmetry, we only need to blow down one of the three NECs: P_1, P_2 or G_0. Thus blowing down P_1 (resp. P_2, or G_0) of $U_1(D_7)$ will give rise to $U_2(A_6)$ (resp. $U_2(D_6)$ and $U_2(A_1 + D_5)$) and the first component of "Exceptional divisor of U_1 to U_m" in the last column is P_1 (resp. P_2, or G_0). When we reach $U_2(A_6)$, we may do the similiar operation as on $U_1(D_7)$ and get $U_3(A_5)$ and $U_3(A_4 + A_1)$.

$No.$	U_1	U_2	U_3	U_4	U_5	U_6	U_7	Exceptional divisor of $U_1 \longrightarrow U_m$
1 2	D_7	A_6	A_5	$A_3 + A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_1 + P_2 + G_2 + G_1 + G_4 + G_5$
				A_4	$A_2 + A_1$	$2A_1$	A_1	$P_1 + P_2 + G_1$ $G_2 + G_4 + G_5$
					A_3	A_2	A_1	$P_1 + P_2 + G_1$ $G_4 + G_5 + G_2$
						$2A_1$	A_1	$P_1 + P_2 + G_1$ $G_4 + G_2 + G_5$
			$A_4 + A_1$	$A_3 + A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_1 + G_0 + P_2 + G_1 + G_4 + G_5$
				A_4	$A_2 + A_1$	$2A_1$	A_1	$P_1 + P_0 + P_2 + G_2 + G_4 + G_5$
					A_3	A_2	A_1	$P_1 + G_2 + P_{-1} + G_7 + G_6 + G_5$
						$2A_1$	A_1	$P_1 + G_0 + P_2 + G_4 + G_2 + G_5$
		D_6	D_5	D_4	A_3	$2A_1$	A_1	$P_2 + G_1 + G_4 + G_5 + G_6 + P_1$
						A_2	A_1	$P_2 + G_1 + G_4 + G_5 + P_1 + G_6$
				A_4	$A_2 + A_1$	$2A_1$	A_1	$P_2 + G_1 + P_1 + G_2 + G_4 + G_5$
					A_3	A_2	A_1	$P_2 + G_1 + P_1 + G_4 + G_5 + G_6$
						$2A_1$	A_1	$P_2 + G_1 + P_1 + G_4 + G_2 + G_6$
		$A_1 + D_5$	D_5	D_4	A_3	A_2	A_1	$G_0 + P_2 + G_4 + G_5 + P_1 + G_6$
						$2A_1$	A_1	$G_0 + P_2 + G_4 + G_5 + G_6 + P_1$
				A_4	$A_2 + A_1$	$2A_1$	A_1	$G_0 + P_2 + P_1 + G_2 + G_4 + G_5$
					A_3	A_2	A_1	$G_0 + P_2 + P_1 + G_4 + G_5 + G_6$
						$2A_1$	A_1	$G_0 + P_2 + P_1 + G_4 + G_2 + G_5$

			A_5	A_3+A_1	A_2+A_1	$2A_1$	A_1	$P_2+P_1+G_2+G_1+G_4+G_5$
				A_4	A_2+A_1	$2A_1$	A_1	$P_2+P_1+G_1+G_2+G_4+G_5$
					A_3	A_2	A_1	$P_2+P_1+G_1+G_4+G_5+G_2$
						$2A_1$	A_1	$P_2+P_1+G_1+G_4+G_2+G_5$
			A_1+A_4	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$G_0+P_1+G_2+P_2+G_4+G_5$
				A_3+A_1	A_2+A_1	$2A_1$	A_1	$G_0+P_1+P_{-1}+G_2+G_7+G_6$
				A_4	A_2+A_1	$2A_1$	A_1	$G_0+P_1+P_2+G_2+G_4+G_5$
					A_3	A_2	A_1	$G_0+P_1+P_2+G_4+G_5+G_2$
						$2A_1$	A_1	$G_0+P_1+P_2+G_4+G_2+G_5$
3	A_3+4A_1	$6A_1$						P_2
4 5	D_5+2A_1	D_4+2A_1	A_3+2A_1	$4A_1$				$P_2+G_1+G_4$
		A_3+3A_1	A_3+2A_1	$4A_1$				$G_0+P_2+G_4$
6	D_4+A_3	$2A_3$	A_3+2A_1	$4A_1$				$P_2+G_1+P_{2,Q}$
		D_4+2A_1	A_3+2A_1	$4A_1$				$P_{2,Q}+P_2+G_1$
		A_3+3A_1	A_3+2A_1	$4A_1$				$G_0+P_2+P_{2,Q}$
7 8	E_7	E_6	D_5	A_4	A_2+A_1	$2A_1$	A_1	$P_1+G_1+G_7+G_6+G_0+G_5$
					A_3	$2A_1$	A_1	$P_1+G_1+G_7+G_0+G_6+G_5$
						A_2	A_1	$P_1+G_1+G_7+G_0+G_5+G_6$
				D_4	A_3	$2A_1$	A_1	$P_1+G_1+G_0+G_5+G_4+G_7$
						A_2	A_1	$P_1+G_1+G_0+G_5+G_7+G_4$
		D_6	D_5	A_4	A_2+A_1	$2A_1$	A_1	$G_0+P_1+G_5+G_4+G_1+G_7$
					A_3	$2A_1$	A_1	$G_0+P_1+G_5+G_1+G_4+G_7$
						A_2	A_1	$G_0+P_1+G_5+G_1+G_7+G_4$
				D_4	A_3	$2A_1$	A_1	$G_0+P_1+G_1+G_5+G_4+G_7$
						A_2	A_1	$G_0+P_1+G_1+G_5+G_7+G_4$

			A_5	A_3+A_1	A_2+A_1	$2A_1$	A_1	$G_0+G_5+G_4+P_1+G_1+G_7$
				A_4	A_2+A_1	$2A_1$	A_1	$G_0+G_5+G_1+G_4+G_1+G_7$
					A_3	$2A_1$	A_1	$G_0+G_5+G_1+G_1+G_4+G_7$
						A_2	A_1	$G_0+G_5+G_1+G_1+G_7+G_4$
9	A_7	A_5+A_1	A_3+2A_1	$4A_1$				$P_1+Q+P_{1,Q}$
		$2A_3$	A_3+2A_1	$4A_1$				$Q+P_1+P_{1,Q}$
10 11	A_7	A_5+A_1	$2A_2+A_1$	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_2+P_1+P_3+G_1+G_3+G_4$
			A_1+A_4	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_2+P_3+P_1+G_1+G_3+G_4$
				A_3+A_1	A_2+A_1	$2A_1$	A_1	$P_2+P_3+G_1+G_2+G_3+G_4$
				A_4	A_2+A_1	$2A_1$	A_1	$P_2+P_3+G_0+P_1+G_5+G_6$
					A_3	A_2	A_1	$P_2+P_3+G_0+G_6+G_5+P_1$
						$2A_1$	A_1	$P_2+P_3+G_0+G_6+P_1+G_5$
		A_2+A_4	$2A_2+A_1$	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_1+P_2+P_3+G_1+G_3+G_4$
			A_4+A_1	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_1+P_3+P_2+G_1+G_3+G_4$
				A_3+A_1	A_2+A_1	$2A_1$	A_1	$P_1+P_3+G_0+P_2+G_6+G_5$
				A_4	A_2+A_1	$2A_1$	A_1	$P_1+P_3+G_1+P_2+G_3+G_4$
					A_3	$2A_1$	A_1	$P_1+P_3+G_1+G_3+G_2+G_4$
						A_2	A_1	$P_1+P_3+G_1+G_3+G_4+G_5$
		A_6	A_4+A_1	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_3+P_1+P_2+G_2+G_3+G_4$
				A_3+A_1	A_2+A_1	$2A_1$	A_1	$P_3+P_1+G_1+G_1+G_3+G_4$
				A_4	A_2+A_1	$2A_1$	A_1	$P_3+P_1+G_1+P_2+G_3+G_4$
					A_3	$2A_1$	A_1	$P_3+P_1+G_1+G_3+P_2+G_4$
						A_2	A_1	$P_3+P_1+G_1+G_3+G_4+G_5$

			A_5	A_3+A_1	A_2+A_1	$2A_1$	A_1	$P_3+G_1+P_2+G_2+G_3+G_4$
				A_4	A_2+A_1	$2A_1$	A_1	$P_3+G_1+G_2+P_2+G_3+G_4$
					A_3	$2A_1$	A_1	$P_3+G_1+G_2+G_3+G_2+G_4$
						A_2	A_1	$P_3+G_1+G_2+G_3+G_4+G_5$
12	$2A_3+A_1$							
13	D_6+A_1	A_5+A_1	A_3+2A_1	$4A_1$				$P_1+G_3+G_0$
14		D_4+2A_1	A_3+2A_1	$4A_1$				$G_0+P_1+G_3$
15 16	A_5+A_2	$3A_2$						P_1
17	D_4+3A_1	$6A_1$						G_0
18 19 20	E_6+A_1	E_6	D_5	A_4	A_2+A_1	$2A_1$	A_1	$P_3+P_1+G_1+G_4+P_2+G_2$
					A_3	$2A_1$	A_1	$P_3+P_1+G_1+P_2+G_4+G_2$
						A_2	A_1	$P_3+P_1+G_1+P_2+G_2+G_4$
				D_4	A_3	$2A_1$	A_1	$P_3+P_1+P_2+G_2+G_5+G_1$
						A_2	A_1	$P_3+P_1+P_2+G_2+G_1+G_5$
		D_5+A_1	D_5	A_4	A_2+A_1	$2A_1$	A_1	$P_2+P_3+G_2+G_5+p_1+G_1$
					A_3	$2A_1$	A_1	$P_2+P_3+G_2+P_1+G_5+G_1$
						A_2	A_1	$P_2+P_3+G_2+P_1+G_1+G_5$
				D_4	A_3	$2A_1$	A_1	$P_2+P_3+P_1+G_2+G_5+G_1$
						A_2	A_1	$P_2+P_3+P_1+G_2+G_1+G_5$
			A_4+A_1	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_2+G_2+G_5+P_3+P_1+G_1$
				A_3+A_1	A_2+A_1	$2A_1$	A_1	$P_2+G_2+G_0+G_3+G_6+G_4$
				A_4	A_2+A_1	$2A_1$	A_1	$P_2+G_2+P_3+G_5+P_1+G_1$
					A_3	$2A_1$	A_1	$P_2++G_2+P_3+P_1+G_5+G_1$
						A_2	A_1	$P_2++G_2+P_3+P_1+G_1+G_5$

		A_5+A_1	$2A_2+A_1$	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$G_0+G_3+P_2+$ $G_2+G_6+G_4$
			A_4+A_1	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$G_0+P_2+G_3+$ $G_2+G_6+G_4$
				A_3+A_1	A_2+A_1	$2A_1$	A_1	$G_0+P_2+G_2+$ $G_3+G_6+G_4$
				A_4	A_2+A_1	$2A_1$	A_1	$G_0+P_2+P_{-1}+$ $G_3+P_1+G_1$
					A_3	A_2	A_1	$G_0+P_2+P_{-1}+$ $P_1+G_1+G_3$
						$2A_1$	A_1	$G_0+P_2+P_{-1}+$ $P_1+G_3+G_1$
21 22	A_3+A_2+ $2A_1$	A_3+3A_1	A_3+2A_1	$4A_1$				$P_4+H_1+P_2$
23 24	$3A_2+A_1$	$3A_2$						P_3
25 26 27	A_6+A_1	A_3+A_2+ A_1	$2A_2+A_1$	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_2+P_4+G_1+$ $G_2+G_4+G_5$
		A_5+A_1	$2A_2+A_1$	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_4+P_2+G_1+$ $G_2+G_4+G_5$
			A_4+A_1	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$P_4+G_2+P_2+$ $G_2+G_4+G_5$
				A_3+A_1	A_2+A_1	$2A_1$	A_1	$P_4+G_1+G_2+$ $G_3+G_4+G_5$
				A_4	A_2+A_1	$2A_1$	A_1	$P_4+G_1+P_1+$ $P_2+P_3+G_6$
					A_3	A_2	A_1	$P_4+G_1+P_1+$ $P_2+G_6+P_2$
						$2A_1$	A_1	$P_4+G_1+P_1+$ $P_3+G_6+P_2$
28 29 30	D_5+A_2	A_3+A_2+ A_1	$2A_2+A_1$	A_2+2A_1	A_2+A_1	$2A_1$	A_1	$G_0+P_3+G_3+$ $G_5+P_1+H_1$
		D_5+A_1	D_5	A_4	A_2+A_1	$2A_1$	A_1	$P_4+H_1+P_3+$ $G_3+P_2+G_1$
					A_3	$2A_1$	A_1	$P_4+H_1+P_3+$ $P_2+G_3+G_1$
						A_2	A_1	$P_4+H_1+P_3+$ $P_2+G_1+G_3$
				D_4	A_3	A_2	A_1	$P_4+H_1+P_2+$ $G_1+P_3+G_4$
						$2A_1$	A_1	$P_4+H_1+P_2+$ $G_1+P_3+G_4$

			$A_4 + A_1$	$A_2 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_4 + P_3 + G_3$ $H_1 + P_2 + G_1$
				$A_3 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_4 + P_3 + P_1$ $G_2 + G_5 + G_4$
				A_4	$A_2 + A_1$	$2A_1$	A_1	$P_4 + P_3 + H_1 +$ $G_3 + P_2 + G_1$
					A_3	$2A_1$	A_1	$P_4 + P_3 + H_1 +$ $P_2 + G_3 + G_1$
						A_2	A_1	$P_4 + P_3 + H_1 +$ $P_2 + G_1 + G_3$
		$A_4 + A_2$	$2A_2 + A_1$	$A_2 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_3 + G_3 + P_4 +$ $H_1 + P_2 + G_1$
			$A_4 + A_1$	$A_2 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_3 + P_4 + G_3 +$ $H_1 + P_2 + G_1$
				$A_3 + A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_4 + P_3 + P_1 +$ $G_2 + G_5 + G_4$
				A_4	$A_2 + A_1$	$2A_1$	A_1	$P_3 + P_4 + H_1 +$ $G_3 + P_2 + G_1$
					A_3	$2A_1$	A_1	$P_3 + P_4 + H_1 +$ $P_2 + G_3 + G_1$
						A_2	A_1	$P_3 + P_4 + H_1 +$ $P_2 + G_1 + G_3$
31	$A_5 + 2A_1$	$A_5 + A_1$	$A_3 + 2A_1$	$4A_1$				$P_3 + P_2 + P_1$
32		$A_3 + 3A_1$	$A_3 + 2A_1$	$4A_1$				$P_2 + P_3 + P_1$
33 34	$A_4 + A_3$	$A_3 + A_2 +$ A_1	$2A_2 + A_1$	$A_2 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_4 + P_5 + H_1 +$ $H_2 + P_3 + G_1$
		$A_4 + A_2$	$2A_2 + A_1$	$A_2 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_5 + P_1 + P_4 +$ $G_2 + P_3 + H_3$
			$A_4 + A_1$	$A_3 + A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_5 + H_1 + P_2 +$ $G_4 + g_3 + G_2$
				$A_2 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_5 + H_1 + P_4 +$ $P_2 + G_3 + G_2$
				A_4	$A_2 + A_1$	$2A_1$	A_1	$P_5 + H_1 + H_2 +$ $P_4 + P_3 + G_1$
					A_3	$2A_1$	A_1	$P_5 + H_1 + H_2$ $P_3 + P_4 + G_1$
						A_2	A_1	$P_5 + H_1 + H_2$ $P_3 + G_1 + G_2$
35 36 37 38	$A_4 + A_2 +$ A_1	$A_3 + A_2 + A_1$	$2A_2 + A_1$	$A_2 + 2A_1$	$A_2 + A_1$	$2A_1$	A_1	$P_6 + G_1 + G_2$ $G_3 + P_4 + H_1$

References

[Artin1] M.Artin: *Some numerical criteria for contractibility of curves on algebraic surfaces*, Amer.J.Math **84** (1962), 485-496.

[Artin2] M.Artin: *On isolated rational singularities of surfaces*, Amer.J.Math **88** (1966), 129-136.

[ATZ] E.Artal Bartolo, H.Tokunaga and D.-Q.Zhang: *Miranda-Persson's Problem on Extremal Elliptic $K3$ Surfaces*, Pacific J.Math., to appear.

[Brenton] L.Brenton: *On Singular complex surfaces with negative canonical bundle, with applications to singular compactifications of C^2 and to 3-dimensional rational singularities*, Math.Ann. **248** (1980), 117-124.

[D] M.Demazure: *Surfaces de del Pezzo*, in: Lecture Notes in Mathematics, Vol. **777**, Springer, 1980.

[DZ] I.Dolgachev and D.-Q. Zhang: *Coble rational surfaces*, Amer.J.Math.**123**(2001), 79-114.

[Fujiki] A.Fujiki, R.Kobayashi, and S.Lu: *On the Fundamental group of Certain Open normal Surfaces*, Saitama Math.J. **11** (1993), 15-20.

[Fu] M.Furushima: *Singular del Pezzo surfaces and analytic compactifications of 3-dimensional complex affine space* $\mathbf{C}^3$, Nagoya Math.J.**104** (1986), 1-28.

[GPZ] R.V.Gurjar, C.R.Pradeep and D.-Q. Zhang: *On Gorenstein Surfaces isomorphic to* $\mathbf{P}^2/G$, preprint, 2000.

[GZ] R.V.Gurjar and D.-Q. Zhang: π_1 *of smooth points of a log del Pezzo surface is finite: I* , J. Math. Sci. Univ. Tokyo **1** (1994), 137-180.

[GZ1] R.V.Gurjar and D.-Q. Zhang: π_1 *of smooth points of a log del Pezzo surface is finite: II* , J. Math. Sci. Univ. Tokyo **2** (1995), 165-196.

[Her] S.Herfurtner : *Elliptic surfaces with four singular fibres*, Math.Ann. **291**(1991), 319-342.

[Hir] U.Schmickler-Hirzebruch : *Elliptische Flächen* über $\mathbf{P}_1\mathbf{C}$ *mit drei Ausnahmefasern und die hypergeometrische Differentialgleichung.* Schriftreihe des Mathematischen Instituts der Universität Münster, 2. Ser. **33**(1985).

[Kas] A.Kas : *Weierstrass normal forms and invariants of elliptic surfaces*, Trans.A.M.S. **225**(1977), 259-266.

[K1] K.Kodaira: *On compact complex analytic surfaces II*, Ann.Math. **77**(1963), 563-626.

[K2] K.Kodaira: *Pluricanonical systems on algebraic surfaces of general type*, J.Math.Soc.Japan Vol.**20**(1968), 170-192.

[KM] S.Keel and J.McKernan: *Rational curves on quasi-projective surfaces*, Memoirs of the A.M.S. **669**(1999).

[KMM] Y.Kawamata, K.Matsuda, K.Matsuki: *Introduction to the Minimal Model Problem*, Advanced Studies in Pure Mathematics **10**(1985), Algebraic Geometry, Sendai.

[M] R.Miranda: *Persson's list of singular fibres for a rational elliptic surface*, Math.Z. **190**(1990), 191-211.

[MP] R.Miranda and U.Persson: *On extremal rational elliptic surfaces*, Math.Z. **193**(1986), 537-558.

[MP1] R.Miranda and U.Persson: *Configurations of I_n fibers on elliptic $K3$ surfaces*, Math.Z. **201**(1989), 339-361.

[MP2] R.Miranda and U.Persson: *Mordell-Weil Groups of extremal elliptic $K3$ surfaces*, Problems in the theory of surfaces and their classification (Cortona, 1988), Symposia Mathematica, XXXII, Academic Press, London,1991, 167-192.

[M1] M.Miyanishi: *Open Algebraic Surfaces*, CRM monograph series, Amer.Math.Soc., 2000.

[MZ1] M.Miyanishi and D.-Q. Zhang : *Gorenstein log del Pezzo surfaces of rank one*, J. of Algebra, **118** (1988), 63-84.

[MZ2] M.Miyanishi and D.-Q. Zhang : *Gorenstein log del Pezzo surfaces,II*, J. of Algebra, **156** (1993), 183-193.

[OS] K.Oguiso and T.Shioda: *The Mordell-Weil Lattice of a Rational Elliptic Surface*, Commentarii Mathematici, Universitatis Sancti Pauli, Vol.**40**(1991), 83-99.

[P] U.Persson : *Configurations of Kordaira fibers on rational elliptic surfaces*, Math.Z.**205** (1990), 1-47.

[Reid] M.Reid : *Chapters on algebraic surfaces*, in "Complex Algebraic Geometry", ed. J. Kollár, IAS/Park City Mathematics Series **3** (1997), 5-159.

[S] T.Shioda: *Elliptic modular surfaces*,J.Math.Soc.Japan **24**(1972),20-59.

[Sh] T.Shioda: *On the Mordell-Weil Lattices*, Commentarii Mathematici, Universitatis Sancti Pauli, Vol.**39**(1990), 211-240.

[SZ] I.Shimada and D.-Q.Zhang : *Classification of extremal elliptic $K3$ surfaces and fundamental groups of open $K3$ surfaces*, Nagoya Math.J.**161**(2001), 23-54.

[T1] S.Takayama : *Simple connectedness of weak Fano varities*, J.Alg.Geom., Vol.**9** (2000), 403-407.

[Ye] Q.Ye: *On extremal elliptic $K3$ surfaces*, J.Korean Math.Soc. **36**(1999), 1091-1113.

[Z1] D.-Q. Zhang : *On Iitaka surfaces*, Osaka J. Math. **24**(1987), 417-460.

[Z2] D.-Q. Zhang : *Logarithmic del Pezzo surfaces of rank one with contractible boundaries*, Osaka J. Math. **25**(1988), no.2, 461-497.

[Z3] D.-Q. Zhang : *Logarithmic del Pezzo surfaces with rational double and triple singular points*, Tôhoku Math. J. (2) **41**(1989), no.3, 399-452.

[Z4] D.-Q. Zhang : *Algebraic surfaces with nef and big anti-canonical divisor* , Math.Proc.Cambridge Philos.Soc. **117**(1995),no.1, 161-163.

[Z5] D.-Q. Zhang : *The fundamental group of the smooth part of a log Fano varity*, Osaka J. Math. **32**(1995), no.3, 637-644.

[Z6] D.-Q. Zhang : *Algebraic surfaces with log canonical singularities and the fundamental groups of their smooth parts*, Trans.Amer.Math.Soc. **348**(1996), no.10, 4175-4184.

Departmant of Mathematics
National University of Singapore
2 Science Drive 2
Singapore 117543
e-mail: mathyeq@hotmail.com